

Publications

6-15-2016

Search for Transient Gravitational Waves in Coincidence With Short-Duration Radio Transients During 2007-2013

B. P. Abbott

California Institute of Technology

K. Gill

Embry-Riddle Aeronautical University, gillk6@my.erau.edu

B. Hughey

Embry-Riddle Aeronautical University, hugheyb@erau.edu

D. Stiles

Embry-Riddle Aeronautical University

M. Szczepańczyk

Embry-Riddle Aeronautical University, szczepam@my.erau.edu

See next page for additional authors

Follow this and additional works at: <https://commons.erau.edu/publication>



Part of the [Physical Sciences and Mathematics Commons](#)

Scholarly Commons Citation

Abbott, B. P., Gill, K., Hughey, B., Stiles, D., Szczepańczyk, M., Zanolin, M., & al., e. (2016). Search for Transient Gravitational Waves in Coincidence With Short-Duration Radio Transients During 2007-2013. *Physical Review D*, 93(12). <https://doi.org/10.1103/PhysRevD.93.122008>

This Article is brought to you for free and open access by Scholarly Commons. It has been accepted for inclusion in Publications by an authorized administrator of Scholarly Commons. For more information, please contact commons@erau.edu.

Authors

B. P. Abbott, K. Gill, B. Hughey, D. Stiles, M. Szczepańczyk, M. Zanolin, and et al.

P. J. Sutton,⁸³ B. L. Swinkels,³⁵ M. J. Szczepański,⁹⁷ M. Tacca,³¹ D. Talukder,⁵⁹ D. B. Tanner,⁵ M. Tápai,⁹⁶ S. P. Tarabrin,⁹ A. Taracchini,³⁰ R. Taylor,¹ T. Theeg,⁹ M. P. Thirugnanasambandam,¹ E. G. Thomas,⁴⁵ M. Thomas,⁶ P. Thomas,³⁸ K. A. Thorne,⁶ E. Thrane,¹¹⁴ S. Tiwari,^{13,89} V. Tiwari,⁸³ K. V. Tokmakov,¹⁰⁵ C. Tomlinson,⁸⁷ M. Tonelli,^{19,20} C. V. Torres^{‡, 86} C. I. Torrie,¹ D. Töyrä,⁴⁵ F. Travasso,^{33,34} G. Traylor,⁶ D. Trifirò,²² M. C. Tringali,^{88,89} L. Trozzo,^{131,20} M. Tse,¹¹ M. Turconi,⁵³ D. Tuyenbayev,⁸⁶ D. Ugolini,¹³² C. S. Unnikrishnan,⁹⁸ A. L. Urban,¹⁷ S. A. Usman,³⁶ H. Vahlbruch,¹⁸ G. Vajente,¹ G. Valdes,⁸⁶ N. van Bakel,¹⁰ M. van Beuzekom,¹⁰ J. F. J. van den Brand,^{62,10} C. Van Den Broeck,¹⁰ D. C. Vander-Hyde,^{36,23} L. van der Schaaf,¹⁰ J. V. van Heijningen,¹⁰ A. A. van Veggel,³⁷ M. Vardaro,^{42,43} S. Vass,¹ M. Vasúth,³⁹ R. Vaulin,¹¹ A. Vecchio,⁴⁵ G. Vedovato,⁴³ J. Veitch,⁴⁵ P. J. Veitch,¹⁰² K. Venkateswara,¹³³ D. Verkindt,⁷ F. Vetrano,^{57,58} A. Viceré,^{57,58} S. Vinciguerra,⁴⁵ D. J. Vine,⁵⁰ J.-Y. Vinet,⁵³ S. Vitale,¹¹ T. Vo,³⁶ H. Vocca,^{33,34} C. Vorwick,³⁸ D. V. Voss,⁵ W. D. Vousden,⁴⁵ S. P. Vyatchanin,⁴⁹ A. R. Wade,²¹ L. E. Wade,¹³⁴ M. Wade,¹³⁴ M. Walker,² L. Wallace,¹ S. Walsh,¹⁷ G. Wang,^{13,58} H. Wang,⁴⁵ M. Wang,⁴⁵ X. Wang,⁷¹ Y. Wang,⁵¹ R. L. Ward,²¹ J. Warner,³⁸ M. Was,⁷ B. Weaver,³⁸ L.-W. Wei,⁵³ M. Weinert,⁹ A. J. Weinstein,¹ R. Weiss,¹¹ T. Welborn,⁶ L. Wen,⁵¹ P. Weßels,⁹ T. Westphal,⁹ K. Wette,⁹ J. T. Whelan,^{113,9} S. E. Whitcomb,¹ D. J. White,⁸⁷ B. F. Whiting,⁵ R. D. Williams,¹ A. R. Williamson,⁸³ J. L. Willis,¹³⁵ B. Willke,^{18,9} M. H. Wimmer,^{9,18} W. Winkler,⁹ C. C. Wipf,¹ H. Wittel,^{9,18} G. Woan,³⁷ J. Worden,³⁸ J. L. Wright,³⁷ G. Wu,⁶ J. Yablon,¹⁰⁷ W. Yam,¹¹ H. Yamamoto,¹ C. C. Yancey,⁶³ M. J. Yap,²¹ H. Yu,¹¹ M. Yvert,⁷ A. Zadrożny,¹¹¹ L. Zangrando,⁴³ M. Zanolini,⁹⁷ J.-P. Zendri,⁴³ M. Zevin,¹⁰⁷ F. Zhang,¹¹ L. Zhang,¹ M. Zhang,¹²⁰ Y. Zhang,¹¹³ C. Zhao,⁵¹ M. Zhou,¹⁰⁷ Z. Zhou,¹⁰⁷ X. J. Zhu,⁵¹ M. E. Zucker,^{1,11} S. E. Zuraw,¹⁰¹ and J. Zweizig¹

(LIGO Scientific Collaboration and Virgo Collaboration)

AND

A. M. Archibald,¹³⁶ S. Banaszak,¹⁷ A. Berndsen,¹³⁷ J. Boyles,¹³⁸ R. F. Cardoso,¹³⁹ P. Chawla,¹⁴⁰ A. Cherry,¹³⁷ L. P. Dartez,⁸⁶ D. Day,¹⁷ C. R. Epstein,¹⁴¹ A. J. Ford,⁸⁶ J. Flanigan,¹⁷ A. Garcia,⁸⁶ J. W. T. Hessels,^{136,142} J. Hinojosa,⁸⁶ F. A. Jenet,⁸⁶ C. Karako-Argaman,¹⁴⁰ V. M. Kaspi,¹⁴⁰ E. F. Keane,¹⁴³ V. I. Kondratiev,^{136,144} M. Kramer,^{145,146} S. Leake,⁸⁶ D. Lorimer,¹³⁹ G. Lunsford,⁸⁶ R. S. Lynch,¹⁴⁷ J. G. Martinez,⁸⁶ A. Mata,⁸⁶ M. A. McLaughlin,¹³⁹ C. A. McPhee,¹³⁷ T. Pennucci,¹⁴⁸ S. Ransom,¹⁴⁹ M. S. E. Roberts,¹⁵⁰ M. D. W. Rohr,¹⁷ I. H. Stairs,¹³⁷ K. Stovall,¹⁵¹ J. van Leeuwen,^{136,142} A. N. Walker,¹⁷ and B. L. Wells^{17,152}

[†]Deceased, May 2015. [‡]Deceased, March 2015.

¹LIGO, California Institute of Technology, Pasadena, CA 91125, USA

²Louisiana State University, Baton Rouge, LA 70803, USA

³Università di Salerno, Fisciano, I-84084 Salerno, Italy

⁴INFN, Sezione di Napoli, Complesso Universitario di Monte S. Angelo, I-80126 Napoli, Italy

⁵University of Florida, Gainesville, FL 32611, USA

⁶LIGO Livingston Observatory, Livingston, LA 70754, USA

⁷Laboratoire d'Annecy-le-Vieux de Physique des Particules (LAPP),

Université Savoie Mont Blanc, CNRS/IN2P3, F-74941 Annecy-le-Vieux, France

⁸University of Sannio at Benevento, I-82100 Benevento,

Italy and INFN, Sezione di Napoli, I-80100 Napoli, Italy

⁹Albert-Einstein-Institut, Max-Planck-Institut für Gravitationsphysik, D-30167 Hannover, Germany

¹⁰Nikhef, Science Park, 1098 XG Amsterdam, The Netherlands

¹¹LIGO, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

¹²Instituto Nacional de Pesquisas Espaciais, 12227-010 São José dos Campos, São Paulo, Brazil

¹³INFN, Gran Sasso Science Institute, I-67100 L'Aquila, Italy

¹⁴INFN, Sezione di Roma Tor Vergata, I-00133 Roma, Italy

¹⁵Inter-University Centre for Astronomy and Astrophysics, Pune 411007, India

¹⁶International Centre for Theoretical Sciences, Tata Institute of Fundamental Research, Bangalore 560012, India

¹⁷University of Wisconsin-Milwaukee, Milwaukee, WI 53201, USA

¹⁸Leibniz Universität Hannover, D-30167 Hannover, Germany

¹⁹Università di Pisa, I-56127 Pisa, Italy

²⁰INFN, Sezione di Pisa, I-56127 Pisa, Italy

²¹Australian National University, Canberra, Australian Capital Territory 0200, Australia

²²The University of Mississippi, University, MS 38677, USA

²³California State University Fullerton, Fullerton, CA 92831, USA

²⁴LAL, Univ. Paris-Sud, CNRS/IN2P3, Université Paris-Saclay, Orsay, France

²⁵Chennai Mathematical Institute, Chennai 603103, India

²⁶Università di Roma Tor Vergata, I-00133 Roma, Italy

²⁷University of Southampton, Southampton SO17 1BJ, United Kingdom

²⁸Universität Hamburg, D-22761 Hamburg, Germany

²⁹INFN, Sezione di Roma, I-00185 Roma, Italy

- ³⁰Albert-Einstein-Institut, Max-Planck-Institut für Gravitationsphysik, D-14476 Potsdam-Golm, Germany
³¹APC, AstroParticule et Cosmologie, Université Paris Diderot, CNRS/IN2P3, CEA/Irfu,
 Observatoire de Paris, Sorbonne Paris Cité, F-75205 Paris Cedex 13, France
³²Montana State University, Bozeman, MT 59717, USA
³³Università di Perugia, I-06123 Perugia, Italy
³⁴INFN, Sezione di Perugia, I-06123 Perugia, Italy
³⁵European Gravitational Observatory (EGO), I-56021 Cascina, Pisa, Italy
³⁶Syracuse University, Syracuse, NY 13244, USA
³⁷SUPA, University of Glasgow, Glasgow G12 8QQ, United Kingdom
³⁸LIGO Hanford Observatory, Richland, WA 99352, USA
³⁹Wigner RCP, RMKI, H-1121 Budapest, Konkoly Thege Miklós út 29-33, Hungary
⁴⁰Columbia University, New York, NY 10027, USA
⁴¹Stanford University, Stanford, CA 94305, USA
⁴²Università di Padova, Dipartimento di Fisica e Astronomia, I-35131 Padova, Italy
⁴³INFN, Sezione di Padova, I-35131 Padova, Italy
⁴⁴CAMK-PAN, 00-716 Warsaw, Poland
⁴⁵University of Birmingham, Birmingham B15 2TT, United Kingdom
⁴⁶Università degli Studi di Genova, I-16146 Genova, Italy
⁴⁷INFN, Sezione di Genova, I-16146 Genova, Italy
⁴⁸RRCAT, Indore MP 452013, India
⁴⁹Faculty of Physics, Lomonosov Moscow State University, Moscow 119991, Russia
⁵⁰SUPA, University of the West of Scotland, Paisley PA1 2BE, United Kingdom
⁵¹University of Western Australia, Crawley, Western Australia 6009, Australia
⁵²Department of Astrophysics/IMAPP, Radboud University Nijmegen, P.O. Box 9010, 6500 GL Nijmegen, The Netherlands
⁵³Artemis, Université Côte d'Azur, CNRS, Observatoire Côte d'Azur, CS 34229, Nice cedex 4, France
⁵⁴MTA Eötvös University, "Lendulet" Astrophysics Research Group, Budapest 1117, Hungary
⁵⁵Institut de Physique de Rennes, CNRS, Université de Rennes 1, F-35042 Rennes, France
⁵⁶Washington State University, Pullman, WA 99164, USA
⁵⁷Università degli Studi di Urbino 'Carlo Bo', I-61029 Urbino, Italy
⁵⁸INFN, Sezione di Firenze, I-50019 Sesto Fiorentino, Firenze, Italy
⁵⁹University of Oregon, Eugene, OR 97403, USA
⁶⁰Laboratoire Kastler Brossel, UPMC-Sorbonne Universités, CNRS,
 ENS-PSL Research University, Collège de France, F-75005 Paris, France
⁶¹Astronomical Observatory Warsaw University, 00-478 Warsaw, Poland
⁶²VU University Amsterdam, 1081 HV Amsterdam, The Netherlands
⁶³University of Maryland, College Park, MD 20742, USA
⁶⁴Center for Relativistic Astrophysics and School of Physics,
 Georgia Institute of Technology, Atlanta, GA 30332, USA
⁶⁵Laboratoire des Matériaux Avancés (LMA), CNRS/IN2P3, F-69622 Villeurbanne, France
⁶⁶Université Claude Bernard Lyon 1, F-69622 Villeurbanne, France
⁶⁷Universitat de les Illes Balears, IAC3—IEEC, E-07122 Palma de Mallorca, Spain
⁶⁸Università di Napoli 'Federico II', Complesso Universitario di Monte S.Angelo, I-80126 Napoli, Italy
⁶⁹NASA/Goddard Space Flight Center, Greenbelt, MD 20771, USA
⁷⁰Canadian Institute for Theoretical Astrophysics, University of Toronto, Toronto, Ontario M5S 3H8, Canada
⁷¹Tsinghua University, Beijing 100084, China
⁷²University of Michigan, Ann Arbor, MI 48109, USA
⁷³National Tsing Hua University, Hsinchu City, 30013 Taiwan, Republic of China
⁷⁴Charles Sturt University, Wagga Wagga, New South Wales 2678, Australia
⁷⁵University of Chicago, Chicago, IL 60637, USA
⁷⁶Caltech CaRT, Pasadena, CA 91125, USA
⁷⁷Korea Institute of Science and Technology Information, Daejeon 305-806, Korea
⁷⁸Carleton College, Northfield, MN 55057, USA
⁷⁹Università di Roma 'La Sapienza', I-00185 Roma, Italy
⁸⁰University of Brussels, Brussels 1050, Belgium
⁸¹Sonoma State University, Rohnert Park, CA 94928, USA
⁸²Texas Tech University, Lubbock, TX 79409, USA
⁸³Cardiff University, Cardiff CF24 3AA, United Kingdom
⁸⁴University of Minnesota, Minneapolis, MN 55455, USA
⁸⁵The University of Melbourne, Parkville, Victoria 3010, Australia
⁸⁶The University of Texas Rio Grande Valley, Brownsville, TX 78520, USA
⁸⁷The University of Sheffield, Sheffield S10 2TN, United Kingdom
⁸⁸Università di Trento, Dipartimento di Fisica, I-38123 Povo, Trento, Italy
⁸⁹INFN, Trento Institute for Fundamental Physics and Applications, I-38123 Povo, Trento, Italy
⁹⁰Montclair State University, Montclair, NJ 07043, USA

- ⁹¹The Pennsylvania State University, University Park, PA 16802, USA
⁹²National Astronomical Observatory of Japan, 2-21-1 Osawa, Mitaka, Tokyo 181-8588, Japan
⁹³School of Mathematics, University of Edinburgh, Edinburgh EH9 3FD, United Kingdom
⁹⁴Indian Institute of Technology, Gandhinagar Ahmedabad Gujarat 382424, India
⁹⁵Institute for Plasma Research, Bhat, Gandhinagar 382428, India
⁹⁶University of Szeged, Dóm tér 9, Szeged 6720, Hungary
⁹⁷Embry-Riddle Aeronautical University, Prescott, AZ 86301, USA
⁹⁸Tata Institute of Fundamental Research, Mumbai 400005, India
⁹⁹INAF, Osservatorio Astronomico di Capodimonte, I-80131, Napoli, Italy
¹⁰⁰American University, Washington, D.C. 20016, USA
¹⁰¹University of Massachusetts-Amherst, Amherst, MA 01003, USA
¹⁰²University of Adelaide, Adelaide, South Australia 5005, Australia
¹⁰³West Virginia University, Morgantown, WV 26506, USA
¹⁰⁴University of Białystok, 15-424 Białystok, Poland
¹⁰⁵SUPA, University of Strathclyde, Glasgow G1 1XQ, United Kingdom
¹⁰⁶IISER-TVM, CET Campus, Trivandrum Kerala 695016, India
¹⁰⁷Northwestern University, Evanston, IL 60208, USA
¹⁰⁸Institute of Applied Physics, Nizhny Novgorod, 603950, Russia
¹⁰⁹Pusan National University, Busan 609-735, Korea
¹¹⁰Hanyang University, Seoul 133-791, Korea
¹¹¹NCBJ, 05-400 Świerk-Otwock, Poland
¹¹²IM-PAN, 00-956 Warsaw, Poland
¹¹³Rochester Institute of Technology, Rochester, NY 14623, USA
¹¹⁴Monash University, Victoria 3800, Australia
¹¹⁵Seoul National University, Seoul 151-742, Korea
¹¹⁶University of Alabama in Huntsville, Huntsville, AL 35899, USA
¹¹⁷ESPCI, CNRS, F-75005 Paris, France
¹¹⁸Università di Camerino, Dipartimento di Fisica, I-62032 Camerino, Italy
¹¹⁹Southern University and A&M College, Baton Rouge, LA 70813, USA
¹²⁰College of William and Mary, Williamsburg, VA 23187, USA
¹²¹Instituto de Física Teórica, University Estadual Paulista/ICTP South American Institute for Fundamental Research, São Paulo SP 01140-070, Brazil
¹²²University of Cambridge, Cambridge CB2 1TN, United Kingdom
¹²³IISER-Kolkata, Mohanpur, West Bengal 741252, India
¹²⁴Rutherford Appleton Laboratory, HSIC, Chilton, Didcot, Oxon OX11 0QX, United Kingdom
¹²⁵Whitman College, 345 Boyer Avenue, Walla Walla, WA 99362 USA
¹²⁶National Institute for Mathematical Sciences, Daejeon 305-390, Korea
¹²⁷Université de Lyon, F-69361 Lyon, France
¹²⁸Hobart and William Smith Colleges, Geneva, NY 14456, USA
¹²⁹Janusz Gil Institute of Astronomy, University of Zielona Góra, 65-265 Zielona Góra, Poland
¹³⁰Andrews University, Berrien Springs, MI 49104, USA
¹³¹Università di Siena, I-53100 Siena, Italy
¹³²Trinity University, San Antonio, TX 78212, USA
¹³³University of Washington, Seattle, WA 98195, USA
¹³⁴Kenyon College, Gambier, OH 43022, USA
¹³⁵Abilene Christian University, Abilene, TX 79699, USA
¹³⁶ASTRON, the Netherlands Institute for Radio Astronomy, Postbus 2, 7990 AA Dwingeloo, The Netherlands
¹³⁷Department of Physics and Astronomy, University of British Columbia, 6224 Agricultural Road, Vancouver, British Columbia V6T 1Z1, Canada
¹³⁸Department of Physics and Astronomy, Western Kentucky University, Bowling Green, KY 42101, USA
¹³⁹Dept. of Physics and Astronomy, West Virginia University, Morgantown, WV 26506, USA
¹⁴⁰Department of Physics & McGill Space Institute, McGill University, 3600 University Street, Montreal, QC H3A 2T8, Canada
¹⁴¹Department of Astronomy, Ohio State University, 140 W. 18th Avenue, Columbus, OH 43210, USA
¹⁴²Anton Pannekoek Institute for Astronomy, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands
¹⁴³SKA Organization, Jodrell Bank Observatory, SK11 9DL, UK
¹⁴⁴Astro Space Center of the Lebedev Physical Institute, Profsoyuznaya str. 84/32, Moscow 117997, Russia
¹⁴⁵Max Planck Institute for Radio Astronomy, Auf dem Hugel 69, 53121 Bonn, Germany
¹⁴⁶The University of Manchester, Oxford Rd, Manchester M13 9PL, UK
¹⁴⁷National Radio Astronomy Observatory, Green Bank, WV 24944, US
¹⁴⁸Department of Astronomy, University of Virginia, Charlottesville, VA 22904-4325, USA
¹⁴⁹National Radio Astronomy Observatory, Charlottesville, VA 22903, USA
¹⁵⁰Eureka Scientific, Inc., 2452 Delmer Street, Suite 100, Oakland, CA 94602-3017, USA

¹⁵¹*Department of Physics and Astronomy, University of New Mexico, Albuquerque, NM, USA*

¹⁵²*Department of Atmospheric Science, Colorado State University, Fort Collins, CO, USA*

We present an archival search for transient gravitational-wave bursts in coincidence with 27 single-pulse triggers from Green Bank Telescope pulsar surveys, using the LIGO, Virgo and GEO interferometer network. We also discuss a check for gravitational-wave signals in coincidence with Parkes fast radio bursts using similar methods. Data analyzed in these searches were collected between 2007 and 2013. Possible sources of emission of both short-duration radio signals and transient gravitational-wave emission include starquakes on neutron stars, binary coalescence of neutron stars, and cosmic string cusps. While no evidence for gravitational-wave emission in coincidence with these radio transients was found, the current analysis serves as a prototype for similar future searches using more sensitive second-generation interferometers.

I. INTRODUCTION

Plausible models for coincident or near-coincident emission of both radio and gravitational-wave (GW) transients exist for a number of astrophysical phenomena, including single neutron stars, merging neutron star binaries and cosmic string cusps. Identification of a GW in close temporal and spatial coincidence with a fast radio burst or other radio transient could place significant constraints on the source of emission, with further constraints possible based on the morphology of the gravitational-wave signal. In this paper we present a search for GWs in coincidence with millisecond-scale duration radio transient pulses. We have conducted externally triggered searches for gravitational waves with the LIGO-Virgo-GEO network in coincidence with both Galactic single-pulse pulsar candidates from the Green Bank Telescope and a sample of cosmological fast radio burst (FRB) candidates from the Parkes Telescope. Individual radio pulses range from 1 to tens of ms in duration and were observed in frequency bands from hundreds of MHz to 1 GHz. The LIGO Scientific Collaboration (LSC) and Virgo Collaboration regularly search for continuous gravitational waves arriving from the direction of known radio pulsars (see, *e.g.* Refs. [2, 3] for recent examples) and a search for gravitational waves in coincidence with a Vela pulsar glitch was conducted previously [10].

The present work marks the first LIGO-Virgo search in coincidence with radio transients and serves as a prototype for searches with advanced interferometers. Given that the origin of these radio transients is currently unclear, our analysis is designed to search broadly for a gravitational-wave transient burst, without requiring a specific type of waveform.

The paper is organized as follows: after briefly describing the network of gravitational-wave interferometers used in this analysis in section II, we discuss possible mechanisms leading to joint emission of \sim few hundred hertz gravitational waves and radio transient signals. We describe the radio data used in the analysis in section IV, followed by the gravitational-wave search methods and results in sections V and VI, respectively. We conclude with a discussion of future prospects for joint analysis of radio and gravitational-wave data in section VII.

II. GRAVITATIONAL-WAVE INTERFEROMETER NETWORK

The LIGO Scientific Collaboration and Virgo Collaboration operate a network of power-recycled Fabry-Perot Michelson interferometers designed to be sensitive to very small relative changes in length (on the order of one part in 10^{21}) of the two orthogonal detector arms. LIGO operates two sites in the United States, one in Livingston Parish, Louisiana and another at the Hanford site in Washington. Both LIGO facilities operate an interferometer with an arm length of 4 km (called L1 and H1, respectively) and Hanford operated an additional smaller, collocated interferometer (H2) until September 2007 [15]. The LIGO Scientific Collaboration also operates a 600 m interferometer, GEO 600, near Hannover, Germany (G1) [28]. The Virgo Collaboration operates a single 3 km interferometer near Cascina, Italy (V1) [17].

Since this paper involves the analysis of radio transients across a period of several years of initial detector data, multiple science runs of these interferometers are used. Data analyzed in this paper are drawn from summer 2007, coincident with LIGO's fifth and Virgo's first science run, as well as late 2009, coincident with LIGO's sixth and Virgo's third science run. FRB candidates discussed in this paper are coincident with GEO 600 Astrowatch data ranging from 2011 to 2013, and in some cases Virgo's fourth science run, which took place in summer 2011. See [4] for a comparison of sensitivities for these instruments from 2007 to 2014.

The LIGO-Virgo network has undergone extensive upgrades to second-generation instruments, and during the first Advanced LIGO observation run made the first direct detection of a gravitational-wave transient [16]. After reaching design sensitivity the Advanced LIGO [7] and Advanced Virgo [1] detectors will have an order of magnitude improvement in range relative to their first-generation counterparts. Additional advanced interferometers are scheduled to join the global network in the future, including Kagra in Japan [59] and a third LIGO site in India [31].

III. POTENTIAL SOURCES OF JOINT RADIO AND GRAVITATIONAL-WAVE EMISSION

There are a number of astrophysical phenomena that may plausibly produce gravitational waves in close coincidence with radio frequency emission. We focus this discussion on a few types of sources which may produce both GWs and radio

pulses with frequency and duration suitable to the instruments being used in this analysis. More detailed discussion can be found in [52].

A. Single neutron stars

Transient gravitational-wave emission can occur when a temporary deformation of a rapidly rotating neutron star creates a quadrupolar moment. Typically, this is believed to happen as a result of crust cracking from magnetic, gravitational or superfluid forces, dubbed a starquake [25, 26], or from other asteroseismic phenomena resulting in the shifting of the neutron star’s crust [35]. Asteroseismology may result in several types of quasinormal oscillatory modes of the neutron star which could produce GW emission. These include torsional modes at low frequencies [72] and the f-mode, with GW emission believed to typically peak around 2 kHz [20]. The amplitude of the GW emission even in optimistic cases, however, is small enough that sensitivity to this type of source will be limited to our own Galaxy even in the advanced detector era.

Radio pulsars result from beamed emission from the poles of a rapidly rotating, highly magnetized neutron star sweeping past the Earth, producing reliably periodic radio signals. The asteroseismic events described above may result in a distinct increase in the rotation rates of these neutron stars, typically followed by a gradual return to their original period. This phenomenon, called a pulsar glitch, has been observed across a large number of pulsars, especially younger ones (see *e.g.* Ref. [44] and references therein). A search for gravitational-wave emission from quasinormal modes in coincidence with the observed glitching of pulsars was the subject of a previous LSC publication [10]. Models for neutron star asteroseismic phenomena similar to those under discussion have also motivated previous gravitational-wave searches in coincidence with soft gamma repeater flares [13].

A related phenomenon to radio pulsars is the rotating radio transient (RRAT). RRATs emit short-duration radio pulses similar in character to pulsars, but are distinguished by their lack of predictable periodic behavior. RRATs may be “dying” pulsars near the end of their life cycles, neutron stars with especially high magnetization, or conventional pulsars of which the observation is often obscured by intervening matter between the pulsar and Earth, although it is also possible that other phenomena may manifest observationally as RRATs [43].

The standard indication of an asteroseismic event in an isolated neutron star is a pulsar glitch, but there are plausible mechanisms that could result in the observation of a transient radio pulse. This could simply be through the pulsar radio emission coming into view from the Earth as the pulsar’s orbit shifts slightly, but there is also some evidence that pulsar-like radio emission can be “switched on” in coincidence with a glitching mechanism [32, 41, 70]. For some models, gravitational waves emitted by neutron stars are predicted to be detectable at a distance scale on the order of kiloparsecs with first generation of interferometers. We therefore consider single neutron stars as possible sources of coincident GW and

radio transient events.

B. Binary neutron star coalescence

The most easily observable transient gravitational-wave signature in the frequency range of LIGO and Virgo is the merger of a binary system of compact objects, specifically neutron stars or black holes. In the final moments before the compact objects merge, the upward sweep in frequency of the gravitational wave emission is predicted to produce a characteristic chirplike signal. Recent evidence suggests that neutron star binary mergers may create at least some fraction of FRBs [34].

Compact binary coalescence is currently the only confirmed source of directly detectable gravitational waves [16]. Once design sensitivity is reached in the advanced detector era, the ground-based network of interferometers is predicted to detect several to a few hundred binary coalescence GW signals per year of operation [8].

There are several models for radio emission in coincidence with a compact binary coalescence GW signal. This may be pulsarlike radio emission, either from the reactivation of the dormant pulsar emission in one of the neutron stars through interactions prior to merger [37] or by a hypermassive neutron star, which may sometimes be produced as an intermediate state before collapse to a black hole [53]. Another possible mechanism is the radiation at radio frequencies as a result of magnetospheric interactions [29].

Given an appropriate density in the surrounding environment, the gravitational waves emitted by a compact binary coalescence may induce electromagnetic radiation through magnetohydrodynamic interactions. While this interaction would directly produce radiation at the same relatively low frequencies as the GWs themselves, upconversion through inverse Compton radiation may result in emission at radio frequencies [46]. This particular magnetohydrodynamic mechanism does not necessarily require neutron star coalescence as the mechanism for production of the GWs, but this class of source is likely to be able to produce GWs of suitable amplitude and may be surrounded by an environment suitable to this mechanism [45].

While the sensitivity of the LIGO/Virgo network to gravitational waves from compact binary coalescence scenarios is dependent on the mass, spin, and other properties of the merging objects, interferometers in the initial detector era were typically sensitive to mergers of two neutron stars out to a distance on the order of 10 Mpc [8].

C. Cosmic strings

Cosmic strings, formed during symmetry breaking in the early Universe, are topological defects thought to be capable of emitting large amounts of energy from their cusps or kinks [21]. A cosmic string cusp may emit gravitational waves with a $f^{-4/3}$ frequency dependence up to a cutoff frequency [24], potentially at frequencies and amplitudes detectable by

ground-based interferometers [5, 58]. The same cusps may produce short-duration linearly polarized radio bursts [23], a mechanism that has previously been proposed as the origin of the original Lorimer burst [68]. Unlike GWs from other sources discussed, cosmic string cusps could theoretically produce detectable GWs at cosmological distances, which makes them particularly interesting in the context of Parkes FRBs with dispersion measures (DMs) indicative of cosmological distances.

D. Other potential sources

The three classes of sources resulting in simultaneous GW and radio emission described above are not an exhaustive list of theoretical joint sources, but most other types of sources are outside the scope of the analyses described in this paper due to the frequency or duration of the predicted GW and/or radio emission not being well suited to the instruments described in this analysis. For example, some scenarios in which gamma-ray bursts (GRBs) may also result in radio emission are not explicitly considered in developing this analysis; prompt radio emission models [57, 67] predict signals at much lower frequencies than the Green Bank and Parkes telescopes can detect, and GRB radio afterglows [47] occur on longer time scales inconsistent with the short radio pulses that are the subject of the searches described in this paper. Core-collapse supernovae have also been proposed as plausible sources of short-duration radio pulses [32] and GW emission. However, we do not explicitly include supernovae among the classes of emission for which we are searching when designing the analysis as there are no observed nearby core-collapse supernovae in close coincidence with the radio transients under consideration.

IV. RADIO PULSE DATA

A. Green Bank single pulse analysis data

The Robert C. Byrd Green Bank Telescope is the world’s largest fully steerable single-dish radio telescope. In the summer of 2007 a drift-scan pulsar survey was conducted in a band of 350 ± 25 MHz [22]. This time frame was during Initial LIGO’s fifth science run and Initial Virgo’s first science run. In addition to the identification of continuously observable pulsars, a “single-pulse” archival search was performed to look for transient emission of millisecond-scale duration radio pulses. The drift-scan team provided LIGO/Virgo with 33 of these observed single-pulse triggers, ten of which were confirmed to originate from sources with repeated emission through followup observations, thus most likely originating from a pulsar or RRAT. Some of the triggers exhibited only a single radio pulse, while others show several pulses within a 2 min window, but in order to be considered a viable astrophysical signal, all pulses were required to exhibit the $1/f^2$ dispersion behavior expected as a result of dispersion in the interstellar medium.

For each of the 33 candidates, right ascension, declination, dispersion measure, and arrival time at solar system barycenter were provided. The dispersion measures provided (between 15 and 170 pc cm $^{-3}$) are in general consistent with a population of sources from within our own Galaxy. For purposes of the gravitational-wave search, barycentric arrival times were adjusted to UTC arrival times at the detector using code previously applied to LIGO pulsar analyses [3] and cross-checked against conversions to the detector frame provided by Green Bank for a subset of triggers.

A survey of the Galaxy’s Northern Celestial Cap was conducted with the Green Bank Telescope in 2009 and 2010 [62]. The single-pulse analysis searching for RRATs or related phenomena was more automated than in the drift-scan analysis and resulted in seven published candidates being reported. These radio triggers corresponded to Initial LIGO’s sixth and Initial Virgo’s third science runs, and were treated identically to drift-scan triggers for GW analysis purposes.

B. Parkes fast radio bursts

The report of FRBs originating from apparently cosmological distances [50, 55, 64] has led to an increased interest in short-duration radio transients. These radio transients resemble the original Lorimer burst [39] reported by Parkes in 2007. Since then Arecibo and Green Bank have each reported an FRB [42, 60]. The origin of these FRBs is unclear, with several possible astrophysical sources and terrestrial backgrounds posited as the origin of these radio bursts. Recent followup observations of the Arecibo burst in particular indicate it is a repeating phenomenon [61], which substantially narrows the range of plausible sources. An observed correlation with another astrophysical signal would do much to clarify the nature of these bursts and help confirm their nature as a previously unknown astrophysical phenomenon.

The FRBs of interest for a gravitational-wave search (i.e. omitting bursts from 2001, prior to the construction of sensitive GW interferometers) have observed fluences ranging from 0.55 to 7.3 Jy ms, observed width from 0.64 to 15.6 ms, and dispersion measures suggesting cosmological origin, ranging from 563 to 1629 pc cm $^{-3}$ [48]. While most of these FRBs did not occur during LIGO/Virgo science runs, we performed a check in the Virgo science run data and GEO 600 Astrowatch data when available.

Given the difference in the dispersion measure and other properties, the FRBs are most likely primarily a distinct class of sources compared to the RRAT-like observations in the previous section. Current evidence points to an astrophysical origin, especially as known perytons exhibit properties distinct from FRBs [51]. Several of the emission mechanisms discussed in Section III are considered viable candidate progenitors, including neutron star and compact binary coalescing systems [18, 36, 65].

C. Short-duration radio transients not analyzed

Potential FRBs from sources other than the above were considered but were not coincident with an active network of GW interferometers. In some cases this was because they occurred during times before sensitive GW data were available [33, 56]. This includes the original Lorimer burst [39]. Neither radio transient reported to be observed in coincidence with a GRB in Ref. [19] was coincident with LIGO/Virgo data [11]. The first observation of Arecibo’s reported FRB [60, 61] occurred during the LIGO and Virgo network upgrade to advanced interferometers and fell during a time in which GEO 600 was not collecting science data as part of its Astrowatch program, with published followup observations also occurring prior to the first Advanced LIGO observation run. Although FRB110523, identified by Green Bank Telescope [42], occurred during Initial Virgo’s fourth science run, Virgo was not taking data at the time of the event.

V. ANALYSIS METHOD

A. Procedure

The search for gravitational waves in coincidence with short radio transients was conducted using the X-PIPELINE analysis package [63]. This software has been used for a number of GW searches in coincidence with astrophysical triggers. The analysis procedure for this search was modeled directly after conceptually similar searches for GWs in coincidence with gamma-ray bursts [6, 11], but the parameters were modified to account for the particular types of GW sources under consideration. Similar adjustments between GRB and radio transient searches can be used to design radio transient searches in advanced interferometers.

For each radio trigger analyzed, we use X-PIPELINE to conduct a coherent search for a GW signal consistent with the location and time of the radio signal. The physical locations of the individual GW interferometers in the network and the antenna patterns based on their orientations are used to reject potential GW signals that are not consistent with the radio signal’s sky location and only signals within ± 2 minutes of the radio trigger are considered. Coherent and incoherent energy combinations are calculated for each potential trigger and a series of two-dimensional cuts is applied to reject triggers physically inconsistent with a GW. The false alarm probability of any surviving triggers after all cuts is estimated based on the “time-lag” method. This utilizes interferometric data outside but near the on-source window, but introduces hundreds of artificial time offsets between the interferometers that are much larger than the time of flight of a real GW signal in order to obtain statistics on the significance of background in which no coherent signal is present. These procedures are discussed in more detail in Ref. [11], with adjustments for our specific analysis as described below.

B. Analysis-specific search parameters

Our frequency range, temporal and spatial coincidence windows, veto methods, and other parameters were selected to handle a range of possible astrophysical emission mechanisms consistent with short radio pulses as discussed in Section III. Where allowed by calibration [9], the frequency range was between 64 Hz and approximately 3 kHz. The majority of GW searches cut off at 2 kHz due to rising shot noise and increased computational costs at higher frequency. However, increasing the upper frequency range for this analysis allows us to include a large subset of possible GW emission from single neutron stars. This increase in frequency also requires us to perform the search over a much denser grid of points on the sky, but the excellent spatial resolution of radio telescopes relative to many other astrophysical observations makes this adjustment feasible.

The on-source time window when searching for GW signals around the radio pulse was taken to be ± 120 s around the observed radio pulse. While it is difficult to exhaustively cover all possible scenarios for time separation between radio and GW emission, the time window selected covers the offsets between emission for the range of scenarios informing the design of the analysis. Since the radio pulse arrival times are corrected for dispersion there is very little additional uncertainty in the time-of-flight difference between the two types of emission. For analyses in coincidence with Green Bank triggers, the angular uncertainty on the sky is taken to be 0.55 deg. This accounts for two effects, including 95% of Green Bank’s total beam width for a 350 MHz signal and including a small adjustment for the drift of the source across the sky during the time span over which X-PIPELINE conducts a test for a self-consistent GW signal.

Unlike recent GW searches in coincidence with GRBs that used similar procedures [6], our background vetoing procedures do not rely on the assumption that the GW signal will be circularly polarized. While this is a reasonable assumption for signals in coincidence with gamma-ray bursts, the greater variety of possible astrophysical sources we consider in this analysis does not justify this assumption.

C. Simulated waveforms

While we do not require a particular signal morphology for our gravitational wave signal, we tune the analysis and characterize its performance based on an ensemble of simulated waveforms. These ten waveforms include:

- damped sinusoids at peak frequencies of 1750 Hz and 2300 Hz and decay time constants of 0.26 s and 0.13 s respectively, representing typical emission expected from neutron star asteroseismic events under different assumptions for the neutron star equation of state [20];
- cosmic string cusp waveforms with upper frequency limits of 300 Hz and 1000 Hz;

- linearly polarized gaussian-envelope sine waves at central frequencies of 235 and 945 Hz with a quality factor of 9;
- circularly polarized gaussian-envelope sine waves at central frequencies of 150 Hz and 300 Hz, also with a quality factor of 9;
- a compact binary coalescence signal from merging 1.4 solar mass neutron stars;
- a compact binary coalescence signal from a 1.4 solar mass neutron star and a 50 solar mass black hole.

These waveforms were selected to broadly represent the types of gravitational wave signals that may occur in coincidence with radio pulses without focusing too heavily on a specific morphology. In addition, the last three bullet points describe specific signals used in previous LIGO searches in order to facilitate sensitivity comparisons with previous work.

VI. ANALYSIS RESULTS

A. Green Bank pulsar surveys

Of the 33 single-pulse radio candidates from the Green Bank drift-scan survey, 25 were analyzable with at least three interferometers in the LIGO-Virgo network. Of the seven RRAT candidates identified in the Northern Celestial Cap survey, only two were analyzable with two or more interferometers in the gravitational wave network.

None of these 27 radio pulses resulted in viable GW candidates. The most significant result for a single candidate was a 2.7% single trial false alarm probability for RRAT 1944-1017, which is completely consistent with background for an ensemble of 27 trials. Table I shows information about each radio candidate, including information about the radio source, as well as GW network and upper limits on h_{rss} (root sum squared strain) for three of the simulated GW waveforms. The last two entries in the table are the Northern Celestial Cap survey triggers.

In addition to individual analysis of the radio candidates, we also perform a weighted binomial test of the p-value distribution of the most significant surviving trigger from the GW analysis, using the same methodology as employed previously in searches for GWs in coincidence with GRBs [4, 11]. This distribution is plotted against expectation in Fig. 1. The test yields a background probability of 30%, which is consistent with the null hypothesis.

Possible association between GRBs and FRBs has been widely discussed (see, *e.g.*, Refs. [19, 27, 34, 38]), with indications that at least a subset of radio bursts may be coupled with gamma-ray bursts. We therefore follow previous LIGO analyses [6] and calculate 90% confidence level exclusion distances, for two of our simulated circularly polarized waveforms, assuming an optimistic standard siren in which $\sim 1\%$ of a solar mass is converted to gravitational-wave energy. A histogram of these distance constraints is shown in Fig. 2. In




FIG. 1. Cumulative distribution of p-values from the analysis of 27 radio triggers from Green Bank for evidence of a GW transient associated with the event. The expected distribution in the absence of a signal is indicated by the dashed line. Points at p-value of unity are triggers with no event in the on-source region after selection cuts.

general, limits in the few to tens of Mpc range indicate that we would be sensitive to a GW signal under these assumptions well outside of our own Galaxy, but at substantially less than the cosmological distances measured for FRBs. For the 150 Hz sine-Gaussian waveform, using standard calculations [14] about $\sim 4 \times 10^{52}$ ergs of energy would have to be emitted for a detectable source emitting isotropically at a distance of 20 Mpc.

B. Parkes Telescope FRBs

We examined a list of 14 FRBs [48] from Parkes, occurring as early as 2001 but primarily concentrated within the last five years. While none of the event times corresponded to science runs for Hanford or Livingston, eight of the FRBs corresponded to times when GEO 600 Astrowatch data were available and two of these also corresponded to data from Initial Virgo's fourth science run. After omitting two of these FRBs for which GEO data were too nonstationary to yield a quality GW analysis, we searched for GWs in coincidence with a total of six Parkes FRBs. Analysis parameters were kept as similar as feasible to the Green Bank drift-scan analysis described previously. However, the upper end of the frequency range was lowered to 1764 Hz due to the range over which GEO data are calibrated [4] and the higher frequency damped sinusoids were left out of the set of GW morphologies simulated. Since the triggers are nominally at cosmological distances and we are unlikely to be sensitive to damped sinusoid-type signals from neutron stars outside our own Galaxy, this limitation is not a major concern.




FIG. 2. Histograms for the sample of Green Bank radio transients of distance exclusions at 90% confidence level for possible GRBs associated with radio transients. Waveforms are circularly polarized sine-Gaussian GW burst models with central frequency of 150 and 300 Hz.

There was no evidence of a gravitational-wave signal for any of these FRBs (the most significant single trial p-value was 0.07), although it should be noted that the smaller GEO interferometer is less sensitive than the larger interferometers. As such we treat this analysis primarily as a check for loud candidates and do not quote sensitivity upper limits for this search. Instead a list of currently published FRBs without evidence for corresponding GW emission is given in Table II. Since these are not consistent in terms of DM or other characteristics with the RRAT-like candidates identified by Green Bank, we do not include these in the binomial test or other distributional studies presented in Figs. 1 and 2.

VII. CONCLUSIONS AND FUTURE PROSPECTS

The searches described in this current paper should be viewed largely as prototypes for future searches with instruments that will eventually be an order of magnitude more sensitive than the best sensitivities presented here. Since much is currently unknown about FRBs and related phenomena, identification of a GW in close coincidence with a radio burst could provide insight into both the distance and, depending on the GW morphology, astrophysical origin of the radio transient.

In addition to more sensitive searches in coincidence with fast radio bursts, efforts are also underway within the LIGO

and Virgo collaborations to analyze radio transients of longer durations resulting from instruments operating at lower frequency than the Green Bank or Parkes telescopes. Since these transients have properties very different than the ones described here and are not generally expected to come from the same sources, substantially different analysis methods are required to address searches for GWs in coincidence with these signals [71].

In the case of fast radio bursts, it is worth noting that arguments based on the Parkes field of view and observation time suggest that if FRBs are in fact of astrophysical origin, the vast majority of FRBs are currently missed by radio telescopes [49, 64]. Accounting for possible anisotropies in the distribution of FRBs and including both mid-galactic and high-latitude survey data, the all-sky rate for FRBs is estimated to be between 1100 to 9600 FRBs per day above a threshold fluence of 4.0 Jy ms [54]. While externally triggered searches can look for signals with amplitudes of as much as ~ 3 lower than all-sky searches [69], if such a population of FRBs generated detectable gravitational-wave signals, statistical arguments suggest they would be likely to show up in all-sky transient searches (e.g. [12]) as well. However, these detections would not be clearly associated with a FRB and thus lack the ability of a multimessenger search in constraining the possible source of FRBs. In the coming years, statistical information on FRBs is likely to be dramatically improved, especially as wider field radio instruments come online [30, 40, 66].

ACKNOWLEDGMENTS

The authors gratefully acknowledge the support of the United States National Science Foundation (NSF) for the construction and operation of the LIGO Laboratory and Advanced LIGO as well as the Science and Technology Facilities Council (STFC) of the United Kingdom, the Max-Planck-Society (MPS), and the State of Niedersachsen/Germany for support of the construction of Advanced LIGO and construction and operation of the GEO600 detector. Additional support for Advanced LIGO was provided by the Australian Research Council. The authors gratefully acknowledge the Italian Istituto Nazionale di Fisica Nucleare (INFN), the French Centre National de la Recherche Scientifique (CNRS) and the Foundation for Fundamental Research on Matter supported by the Netherlands Organisation for Scientific Research, for the construction and operation of the Virgo detector and the creation and support of the EGO consortium. The authors also gratefully acknowledge research support from these agencies as well as by the Council of Scientific and Industrial Research of India, Department of Science and Technology, India, Science & Engineering Research Board (SERB), India, Ministry of Human Resource Development, India, the Spanish Ministerio de Economía y Competitividad, the Conselleria d'Economia i Competitivitat and Conselleria d'Educació, Cultura i Universitats of the Govern de les Illes Balears, the National Science Centre of Poland, the European Commission, the Royal Society, the Scottish Funding Council, the Scottish Universities Physics Alliance, the Hungarian Scien-

-
- [1] <https://wwwcascina.virgo.infn.it/advirgo/docs.html>.
- [2] Aasi, J. *et al.*, Phys. Rev. D **88**, 102002 (2013).
- [3] Aasi, J. *et al.*, Astrophys. J. **785**, 119 (2014).
- [4] Aasi, J. *et al.*, Phys. Rev. D **89**, 122004 (2014).
- [5] Aasi, J. *et al.*, Phys. Rev. Lett. **112**, 131101 (2014).
- [6] Aasi, J. *et al.*, Phys. Rev. Lett. **113**, 011102 (2014).
- [7] Aasi, J. *et al.*, Class. Quant. Grav. **32**, 074001 (2015).
- [8] Abadie, J. *et al.*, Class. Quant. Grav. **27**, 173001 (2010).
- [9] Abadie, J. *et al.*, Nucl. Inst. Meth. Phys. Res A**624**, 223 (2010).
- [10] Abadie, J. *et al.*, Phys. Rev. D **83**, 042001 (2011).
- [11] Abadie, J. *et al.*, Astrophys. J. **760**, 12 (2012).
- [12] Abadie, J. *et al.*, Phys. Rev. D **85**, 122007 (2012).
- [13] Abbott, B. P. *et al.*, Phys. Rev. Lett. **101**, 211102 (2008).
- [14] Abbott, B. P. *et al.*, Phys. Rev. D **77**, 062004 (2008).
- [15] Abbott, B. P. *et al.*, Reports on Progress in Physics **72**, 076901 (2009).
- [16] Abbott, B. P. *et al.*, Phys. Rev. Lett. **116**, 061102 (2016).
- [17] Accadia, T. *et al.*, Journal of Instrumentation **7**, P03012 (2012).
- [18] Bannister, K. W. and Madsen, G. J., Mon. Not. R. Astron. Soc. **440**, 353 (2014).
- [19] Bannister, K. W., Murphy, T., Gaensler, B. M., and Reynolds, J. E., Astrophys. J. **757**, 38 (2012).
- [20] Benhar, O., Ferrari, V., and Gualtieri, L., Phys. Rev. D **70**, 124015 (2004).
- [21] Berezinsky, V., Hnatyk, B., and Vilenkin, A., Baltic Astronomy **13**, 289 (2004).
- [22] Boyles, J. *et al.*, Astrophys. J. **763**, 80 (2013).
- [23] Cai, Y. F., Sabancilar, E., and Vachaspati, T., Phys. Rev. D **85**, 023530 (2012).
- [24] Damour, T. and Vilenkin, A., Phys. Rev. Lett. **85**, 3761 (2000).
- [25] Duncan, R. C., Astrophys. J. Lett. **498**, L45 (1998).
- [26] Franco, L. M., Link, B., and Epstein, R. I., Astrophys. J. **543**, 987 (2000).
- [27] Gao, H., Li, Z., and Zhang, B., Astrophys. J. **788**, 189 (2014).
- [28] Grote, H. *et al.*, Class. Quant. Grav. **27**, 084003 (2010).
- [29] Hansen, B. M. S. and Lyutikov, M., Mon. Not. R. Astron. Soc. **322**, 695 (2001).
- [30] Hassall, T. E., Keane, E. F., and Fender, R. P., Mon. Not. R. Astron. Soc. **436**, 371 (2013).
- [31] Iyer, B. *et al.*, LIGO-India Tech. rep. <https://dcc.ligo.org/cgi-bin/DocDB/ShowDocument?docid=75988> (2011).
- [32] Keane, E. F., Kramer, M., Lyne, A. G., Stappers, B. W., and McLaughlin, M. A., Mon. Not. R. Astron. Soc. **415**, 3065 (2011).
- [33] Keane, E. F., Stappers, B. W., Kramer, M., and Lyne, A. G., Mon. Not. R. Astron. Soc. **425**, L71 (2012).
- [34] Keane, E. F. *et al.*, Nature **530**, 453 (2016).
- [35] Kokkotas, K. D. and Schmidt, B. G., Living Rev. Rel. **2**, 2 (1999).
- [36] Li, X., Zhou, B., He, H.-N., Fan, Y.-Z., and Wei, D. M., Astrophys. J. **797**, 33 (2014).
- [37] Lipunov, V. M. and Panchenko, I. E., Astronomy and Astrophysics **312**, 937 (1996).
- [38] Lipunov, V. M. and Pruzhinskaya, M. V., Mon. Not. R. Astron. Soc. **440**, 1193 (2014).
- [39] Lorimer, D. R., Bailes, M., McLaughlin, M. A., Narkevic, D. J., and Crawford, F., Science **318**, 777 (2007).
- [40] Lorimer, D. R., Karastergiou, A., McLaughlin, M. A., and Johnston, S., Mon. Not. R. Astron. Soc. **436**, L5 (2013).
- [41] Lyne, A. G. *et al.*, Mon. Not. R. Astron. Soc. **400**, 1439 (2009).
- [42] Masui, K. *et al.*, Nature **528**, 523 (2015).
- [43] McLaughlin, M. A. *et al.*, Nature **439**, 817 (2006).
- [44] Middleitch, J., Marshall, F. E., Wang, Q. D., Gotthelf, E. V., and Zhang, W., Astrophys. J. **652**, 1531 (2006).
- [45] Moortgat, J. and Kuijpers, J., Proceedings of the XXII Texas Symposium on Relativistic Astrophysics (Stanford University, 2004), PSN 1230.
- [46] Moortgat, J. and Kuijpers, J., Phys. Rev. D **70**, 023001 (2004).
- [47] Nakar, E., Phys. Rep. **442**, 166 (2007).
- [48] Petroff, E. *et al.*, arXiv:1601.03547.
- [49] Petroff, E. *et al.*, Astrophys. J. **789**, L26 (2014).
- [50] Petroff, E. *et al.*, Mon. Not. R. Astron. Soc. **447**, 246 (2015).
- [51] Petroff, E. *et al.*, Mon. Not. R. Astron. Soc. **451**, 3933 (2015).
- [52] Predoi, V. *et al.*, Class. Quant. Grav. **27**, 084018 (2010).
- [53] Pshirkov, M. and Postnov, K., Astrophysics and Space Science **330**, 13 (2010).
- [54] Rane, A. *et al.*, Mon. Not. R. Astron. Soc. **455**, 2207 (2016).
- [55] Ravi, V., Shannon, R., and Jameson, A., Astrophys. J. Lett. **799**, L5 (2015).
- [56] Rubio-Herrera, E., Stappers, B. W., Hessels, J. W. T., and Braun, R., Mon. Not. R. Astron. Soc. **428**, 2857 (2013).
- [57] Sagiv, A. and Waxman, E., Astrophys. J. **574**, 861 (2002).
- [58] Siemens, X., Creighton, J., Maor, I., Majumder, S. R., Cannon, K., and Read, J., Phys. Rev. D **73**, 105001 (2006).
- [59] Somiya, K. *et al.*, Class. Quant. Grav. **29**, 124007 (2012).
- [60] Spitler, L. G. *et al.*, Astrophys. J. **790**, 101 (2014).
- [61] Spitler, L. G. *et al.*, Nature **531**, 202 (2016).
- [62] Stovall, K. *et al.*, Astrophys. J. **791**, 67 (2014).
- [63] Sutton, P. J. *et al.*, New Journal of Physics **12**, 053034 (2010).
- [64] Thornton, D. *et al.*, Science **341**, 53 (2013).
- [65] Totani, T., Publ. Astron. Soc. Jpn. **65**, L12 (2013).
- [66] Trott, C. M., Tingay, S. J., and Wayth, R. B., Astrophys. J. Lett. **776**, L16 (2013).
- [67] Usov, V. V. and Katz, J. I., Astronomy and Astrophysics **364**, 655 (2000).
- [68] Vachaspati, T., Phys. Rev. Lett. **101**, 141301 (2008).
- [69] Was, M., Ph.D. thesis, Universite Paris XI (2011).
- [70] Weltevrede, P., Johnston, S., and Espinoza, C. M., AIP Conference Proceedings **1357**, 109 (2011).
- [71] Yancey, C. *et al.*, Astrophys. J. **812**, 168 (2015).
- [72] Zink, B., Lasky, P. D., and Kokkotas, K. D., Phys. Rev. D **85**, 024030 (2012).