

Publications

5-10-2006

The Mystery Deepens: Spitzer Observations of Cool White Dwarfs

Mukremin Kilic

University of Texas at Austin, kilic@ou.edu

Ted von Hippel

University of Texas at Austin, vonhippt@erau.edu

et al.

Follow this and additional works at: <https://commons.erau.edu/publication>



Part of the [Stars, Interstellar Medium and the Galaxy Commons](#)

Scholarly Commons Citation

Kilic, M., von Hippel, T., & al., e. (2006). The Mystery Deepens: Spitzer Observations of Cool White Dwarfs. *The Astrophysical Journal*, 656(2). Retrieved from <https://commons.erau.edu/publication/264>

This Article is brought to you for free and open access by Scholarly Commons. It has been accepted for inclusion in Publications by an authorized administrator of Scholarly Commons. For more information, please contact commons@erau.edu.

THE MYSTERY DEEPENS: *SPITZER* OBSERVATIONS OF COOL WHITE DWARFS

MUKREMIN KILIC,¹ TED VON HIPPEL,¹ FERGAL MULLALLY,¹ WILLIAM T. REACH,² MARC J. KUCHNER,³
D. E. WINGET,¹ AND ADAM BURROWS⁴

Received 2005 October 14; accepted 2006 January 4

ABSTRACT

We present 4.5 and 8 μm photometric observations of 18 cool white dwarfs obtained with the *Spitzer Space Telescope*. Our observations demonstrate that four white dwarfs with $T_{\text{eff}} < 6000$ K show slightly depressed mid-infrared fluxes relative to white dwarf models. In addition, another white dwarf with a peculiar optical and near-infrared spectral energy distribution (LHS 1126) is found to display significant flux deficits in *Spitzer* observations. These mid-infrared flux deficits are not predicted by the current white dwarf models including collision-induced absorption due to molecular hydrogen. We postulate that either the collision-induced absorption calculations are incomplete or there are other unrecognized physical processes occurring in cool white dwarf atmospheres. The spectral energy distribution of LHS 1126 surprisingly fits a Rayleigh-Jeans spectrum in the infrared, mimicking a hot white dwarf with effective temperature well in excess of 10^5 K. This implies that the source of this flux deficit is probably not molecular absorption but some other process.

Subject headings: infrared: stars — stars: individual (LHS 1126, WD 0038–226) — white dwarfs

Online material: color figure

1. INTRODUCTION

The Sloan Digital Sky Survey (Adelman-McCarthy et al. 2006) has increased the number of known field cool white dwarfs from tens of objects to thousands (Kilic et al. 2006; Harris et al. 2006). In addition, *Hubble Space Telescope* observations of the globular clusters M4 (Hansen et al. 2004) and ω Cen (Monelli et al. 2005) resulted in the discovery of more than two thousand white dwarfs. A careful analysis of these large data sets require a complete understanding of the structure and evolution of the white dwarf stars. The main uncertainty in the white dwarf luminosity functions derived from these data sets is caused by the inadequate description of energy transport in white dwarf atmospheres, which affects both the cooling rate and observational appearance of these stars (Hansen 1998).

Cool white dwarfs have atmospheres dominated by hydrogen or helium. Both hydrogen and helium are neutral below 5000 K, and the primary opacity source in H-rich cool ($T_{\text{eff}} \leq 5500$ K) white dwarf atmospheres is believed to be collision-induced absorption (CIA; Frommhold 1993) of molecular hydrogen (Bergeron et al. 1995; Hansen 1998; Saumon & Jacobson 1999). H-rich white dwarfs are predicted to become redder as they cool until the effects of CIA become significant below 5500 K. CIA opacity is strongly wavelength dependent and is expected to produce broad absorption features in the near-infrared. This flux deficiency can be seen in several stars, designated as ultracool white dwarfs (Oppenheimer et al. 2001; Harris et al. 2001; Gates et al. 2004; Farihi 2004, 2005).

There are two known opacity mechanisms in pure He-rich white dwarf atmospheres: Rayleigh scattering and He⁻ free-free

absorption. Rayleigh scattering is thought to be the dominant opacity source, and therefore He-rich white dwarfs are expected to have blackbody-like spectral energy distributions (SEDs). On the other hand, Kowalski et al. (2005) argued that He⁻ free-free absorption may be 2–3 orders of magnitude more significant, which would make it the dominant opacity source. In addition, Kowalski & Saumon (2004) showed that the atmosphere of He-rich white dwarfs with $T_{\text{eff}} < 10000$ K should be treated as a dense fluid rather than an ideal gas and that refraction effects become important. The density can be as high as 2 g cm^{-3} in these atmospheres, and the index of refraction departs significantly from unity. Kowalski et al. (2005) used their updated models to match the observed sequence of cool white dwarfs from Bergeron et al. (1997, hereafter BRL97) and suggested that the coolest white dwarfs have mixed H/He atmospheres.

Bergeron & Leggett (2002) tried to fit the optical and near-infrared photometry of two ultracool white dwarfs, LHS 3250 and SDSS 1337+00, and ruled out their models for pure hydrogen atmospheric composition for these stars. They found that the overall SEDs of these stars can be better fitted with mixed H/He models, yet the peak of the SEDs near 6000 Å is predicted to be too narrow. There are only three white dwarfs with significant CIA that fit the current white dwarf models in the optical and near-infrared. WD 0346+246 (Oppenheimer et al. 2001) and GD 392B (Farihi 2004) both require low-gravity ($\log g \sim 7$), mixed H/He model atmosphere solutions. In addition, Bergeron et al. (1994) found a 5400 K, $\log g = 7.9$, and $\log N(\text{He})/N(\text{H}) = 0.8$ model atmosphere solution for LHS 1126. BRL97 used new CIA opacity calculations and suggested that the helium abundance in this white dwarf is $\log N(\text{He})/N(\text{H}) = 1.86$.

The *Spitzer Space Telescope* (Werner et al. 2004) opened a new window into the universe by enabling accurate mid-infrared photometry of faint objects (microjansky-level sensitivity). In order to understand the CIA opacity, and other unrecognized sources of opacity in cool white dwarf atmospheres, we used *Spitzer* to observe nearby, relatively bright, cool white dwarfs. In this paper, we present our mid-infrared photometry for 18 cool white dwarfs including LHS 1126. Our observations are

¹ Department of Astronomy, University of Texas at Austin, 1 University Station C1400, Austin, TX 78712; kilic@astro.as.utexas.edu.

² *Spitzer* Science Center, MS 220-6, California Institute of Technology, Pasadena, CA 91125.

³ NASA Goddard Space Flight Center, Greenbelt, MD 20771.

⁴ Department of Astronomy, and Steward Observatory, University of Arizona, 933 North Cherry Avenue, Tucson, AZ 85721.

TABLE 1
INFRARED PHOTOMETRY OF COOL WHITE DWARFS

Object	Spectral Type	T_{eff} (K) ^a	F_J (mJy)	F_H (mJy)	F_K (mJy)	$F_{4.5 \mu\text{m}}$ (mJy)	$F_{8 \mu\text{m}}$ (mJy)
WD 0009+501	DAP	6540/6683	6.40 ± 0.12	5.14 ± 0.10	3.53 ± 0.07	0.94 ± 0.05	0.32 ± 0.02
WD 0018–267	DA	.. /5498	15.88 ± 0.29	14.72 ± 0.30	10.51 ± 0.20	2.79 ± 0.09	1.02 ± 0.04
WD 0038–226	C ₂ H:	5400/...	7.34 ± 0.14	4.14 ± 0.09	2.13 ± 0.05	0.48 ± 0.02	0.19 ± 0.03
WD 0101+048 ^b	DA	8080/8160	6.32 ± 0.12	4.49 ± 0.09	2.86 ± 0.06	0.75 ± 0.03	0.29 ± 0.03
WD 0126+101 ^b	DA	8500/8700	3.89 ± 0.07	2.69 ± 0.06	1.73 ± 0.04	0.45 ± 0.02	0.16 ± 0.07
WD 0141–675	DA	.. /6469	11.37 ± 0.20	8.85 ± 0.18	6.20 ± 0.12	1.64 ± 0.06	0.59 ± 0.03
WD 0552–041	DZ	5060/5016	9.63 ± 0.18	7.35 ± 0.15	5.17 ± 0.10	1.80 ± 0.06	0.72 ± 0.07
WD 0553+053	DAP	5790/5853	10.73 ± 0.19	8.36 ± 0.17	5.79 ± 0.11	1.64 ± 0.05	0.51 ± 0.04
WD 0752–676	DA	5730/5774	12.94 ± 0.23	10.47 ± 0.21	7.57 ± 0.15	2.06 ± 0.08	0.77 ± 0.04
WD 0839–327 ^b	DA	.. /8978	37.26 ± 0.68	24.81 ± 0.51	16.04 ± 0.32	4.01 ± 0.14	1.45 ± 0.06
WD 0912+536	DCP	7160/7273	7.57 ± 0.14	5.32 ± 0.11	3.72 ± 0.07	1.10 ± 0.04	0.41 ± 0.03
WD 1055–072	DC	7420/7252	4.95 ± 0.09	3.45 ± 0.07	2.69 ± 0.06	0.64 ± 0.02	0.25 ± 0.03
WD 1121+216 ^c	DA	7490/7540	5.93 ± 0.11	4.39 ± 0.09	2.91 ± 0.06	0.81 ± 0.03	0.34 ± ...
WD 1223–659	DA	.. /7793	7.39 ± 0.14	5.10 ± 0.12	3.20 ± 0.07	0.90 ± 0.04	0.36 ± 0.07
WD 1748+708	DXP	5590/5964	13.15 ± 0.24	9.98 ± 0.20	6.62 ± 0.13	1.89 ± 0.06	0.70 ± 0.04
WD 1756+827	DA	7270/7285	5.61 ± 0.10	4.18 ± 0.09	2.83 ± 0.06	0.75 ± 0.02	0.26 ± 0.04
WD 1953–011	DAP	7920/7851	9.43 ± 0.17	6.29 ± 0.13	4.15 ± 0.09	1.13 ± 0.04	0.41 ± 0.02
WD 2140+207	DQ	8860/9407	10.24 ± 0.18	6.90 ± 0.14	4.52 ± 0.09	1.19 ± 0.05	0.41 ± 0.03

^a Estimated T_{eff} from BLR01 vs. this study.

^b Suspected or known double degenerate.

^c The 8 μm photometry of WD 1121+216 is affected by a nearby star.

discussed in § 2, while an analysis of these data and results is discussed in § 3.

2. TARGET SELECTION AND OBSERVATIONS

We selected our targets from the spectroscopically confirmed white dwarf catalog (McCook & Sion 1999) and required that they have $K_S \leq 15$ mag as measured by the Two Micron All Sky Survey (2MASS).⁵ Observations reported here were obtained as part of the *Spitzer* Cycle 1 GO-Program 2313 (P.I.: Marc Kuchner). We used the Infrared Array Camera (IRAC; Fazio et al. 2004) to obtain 4.5 and 8 μm (effective wavelengths of 4.493 and 7.782 μm) photometry of 18 cool white dwarfs with $5000 \text{ K} < T_{\text{eff}} < 9000 \text{ K}$. An integration time of 30 s per dither, with five dithers for each target, was used (150 s total integration time).

We used the products of the *Spitzer* Science Center pipeline, the Basic Calibrated Data (BCD) frames and the Post-BCD frames (mosaics), for our analysis. We used the IRAF⁶ phot routine to perform aperture photometry on individual BCD frames. Experience showed that the point response function for the IRAC instrument is not well defined, and we obtained better results with aperture photometry than with point-spread-function-fitting photometry. In order to maximize the signal-to-noise ratio, we used a 5 pixel aperture for bright, isolated objects and 2 or 3 pixel apertures for faint objects or objects in crowded fields. We corrected the resulting fluxes by the aperture correction factors determined by the IRAC team (see the IRAC Data Handbook). We checked the results from 2, 3, and 5 pixel apertures for each object and found them to be consistent within the errors.

Following the IRAC calibration procedure, corrections for the location of the source in the array were taken into account

before averaging the fluxes of each of the five frames at each wavelength. We also performed photometry on the mosaic images and found the results to be consistent with the photometry from individual frames. We divided the estimated fluxes by the color corrections for a Rayleigh-Jeans spectrum (Reach et al. 2005b). These corrections are 1.0121 and 1.0339 for the 4.5 and 8 μm bands, respectively. Based on the work of Reach et al. (2005b), we expect that our IRAC photometry is calibrated to an accuracy of 3%. The average fluxes measured from the *Spitzer* images along with the 2MASS photometry, spectral types, and temperatures for our objects are given in Table 1. The error bars were estimated from the observed scatter in the five images (corresponding to five dither positions) plus the 3% absolute calibration error, added in quadrature.

3. RESULTS

Figure 1 shows the optical and infrared SEDs (*triangles*) of the 18 cool white dwarfs (ordered in T_{eff}) that we observed with the IRAC instrument. Most of our objects were observed by Bergeron et al. (2001; hereafter BLR01) and therefore have accurate *BVRI* photometry and temperature determinations. For the four DA white dwarfs not included in BLR01's analysis (WD 0018–267, WD 0141–675, WD 0839–327, and WD 1223–659), we used the *UBV* photometry from the McCook & Sion (1999) catalog. The near-infrared photometry comes from the 2MASS Point Source Catalog.

The expected fluxes from synthetic photometry of white dwarf model atmospheres (integrated over the filter bandpasses; kindly made available to us by D. Saumon and D. Koester) are shown as circles in Figure 1. These models include CIA opacities for H-rich objects with $T_{\text{eff}} < 7000 \text{ K}$. We normalized the model atmospheres to the observed SEDs in the *V* band (0.55 μm), with the exception of WD 1748+708 (G240-72). WD 1748+708 shows a 15% deep, 2000 \AA wide absorption feature centered at $\sim 5300 \text{ \AA}$ that affects its *BVR* photometry (BRL97). Hence, we matched its model to the observations in the *I* band. Solid lines represent the same models normalized to the observed SEDs

⁵ VizieR Online Data Catalog, 2246 (R. M. Cutri et al., 2003)

⁶ IRAF is distributed by the National Optical Astronomy Observatory, which is operated by the Association of Universities for Research in Astronomy (AURA), Inc., under cooperative agreement with the National Science Foundation (NSF).




FIG. 1.—SEDs of cool white dwarfs observed with the *Spitzer Space Telescope*. The observed fluxes are shown as triangles with error bars, whereas the expected flux distributions from synthetic photometry of white dwarf models (kindly made available to us by D. Saumon and D. Koester) are shown as circles. Solid lines represent the same models normalized to the observed SEDs in the *J* band. The object name, atmospheric composition, T_{eff} , and spectral type are given in each panel.

in the *J* band. Changing the normalization of the models to the *J* band does not change our results significantly since the models and observations agree fairly well between the *V* and *J* bands.

In order to derive new T_{eff} values for our objects, we used pure H and pure He white dwarf models ($\log g = 8$) and employed a χ^2 minimization technique to fit *V*, *RI* (if available), *JHK*, and 4.5 and 8 μm photometry. The *V* and *R* photometry was omitted in our analysis of WD 1748+708 because of the strong absorption feature observed in these filters. Our estimated T_{eff} values for the eight DA white dwarfs that are in common with the BLR01 sample are on average slightly hotter ($+66 \pm 81$ K) than the BLR01 temperatures, although they are consistent within the errors.

3.1. White Dwarfs with $T_{\text{eff}} < 7000$ K

A comparison of observed versus expected SEDs from white dwarf model atmospheres shows that all of the cool white dwarfs with $T_{\text{eff}} < 6000$ K (except the DZ white dwarf WD 0552–041) show slightly depressed mid-infrared fluxes relative to white dwarf models. In addition, WD 0038–226 (LHS 1126) displays significant flux deficits (more than 5σ in each band) in the mid-infrared. There are also two more stars with $6000 \text{ K} < T_{\text{eff}} < 7000$ K (WD 0009+501 and WD 0141–675) that display $\sim 1\sigma$ flux deficits. Figures 2 and 3 show the observed flux

ratios in different filters for the white dwarfs in our sample. H- and He-rich white dwarfs are shown as filled and open circles, respectively. White dwarfs with uncertain spectral types (LHS 1126 and WD 1748+708) are shown as triangles. Cooling tracks for H- (solid line) and He-rich (dashed line) white dwarfs, along with a blackbody cooling track (dotted line) are also shown. Figure 2 shows that H-rich white dwarfs with $T_{\text{eff}} < 7000$ K emit less flux at 8 μm than predicted from the white dwarf models. In addition, their infrared colors (Fig. 3) require most of them to be warmer than 6500 K; they mimic warmer/bluer objects in the infrared.

LHS 1126 shows strong molecular features in the optical that are thought to be due to C_2H . These absorption features are blueshifted by about 150 \AA compared to the C_2 Swan bands. Several investigators tried to explain these features in terms of either pressure shifts in a He-dominated atmosphere or magnetic displacements of the Swan bands (see Schmidt et al. 1995). However, Bergeron et al. (1994) and Schmidt et al. (1995) showed that both scenarios failed for this star. In addition, Schmidt et al. (1995) suggested that C_2H is the most probable molecule to form under the conditions in LHS 1126. WD 1748+708 shows a very broad feature in the optical (see the discussion above) and has a ~ 200 MG magnetic field (Angel 1977). BLR01 suggested that this broad feature may be explained as a C_2H molecular




FIG. 1.—Continued




FIG. 2.—Ratio of observed fluxes in V , J , and $8 \mu\text{m}$ bands for DA (filled circles) and DB (open circles) white dwarfs. White dwarfs with uncertain spectral types are shown as triangles. Expected cooling tracks for DA (5000–60000 K, solid line) and DB white dwarfs (5000–8000 K, dashed line) are also shown. A dotted line shows the expected sequence for a blackbody (5000– 10^5 K).




FIG. 3.—Ratio of observed fluxes in J , H , and $4.5 \mu\text{m}$ bands. The symbols are the same as in Fig. 2. Note that the discontinuity at 7000 K is due to transition from Koester models to Saumon models.

feature broadened by the strong magnetic field. Hence, all of the stars with mid-infrared flux deficits have either H-rich atmospheres or show trace amounts of hydrogen.

The coolest white dwarf in our sample is a DZ white dwarf, WD 0552–041, with an estimated $T_{\text{eff}} = 5016$ K. The observed SED fits the model predictions fairly well for this star with the exception of the B photometry, which is probably affected by the metals in the photosphere (Wolff et al. 2002). The origin of metals in white dwarf atmospheres has been a mystery for a long time (Zuckerman et al. 2003). The discovery of debris disks around the white dwarfs G29–38 (Zuckerman & Becklin 1987; Reach et al. 2005a) and GD 362 (Becklin et al. 2005; Kilic et al. 2005) suggests that accretion from a debris disk may explain the observed metal abundances in DAZ white dwarfs. Nevertheless, we do not see any mid-infrared excess around the cool DZ white dwarf WD 0552–041.

3.2. White Dwarfs with $T_{\text{eff}} > 7000$ K

The remaining 10 objects with $T_{\text{eff}} > 7000$ K do not show any excess or deficit in their flux distributions. There are three known or suspected double degenerate stars in our sample (WD 0101+048, WD 0126+101, and WD 0839–327; Maxted et al. 2000). We do not see any evidence of a composite SED for these objects; if they are equal-mass binaries, the primary and secondary stars must have similar temperatures. Limits on possible substellar companions or debris disks around these stars, as well as ~ 100 other white dwarfs, will be discussed in a future publication (T. von Hippel et al. 2006, in preparation). One caveat seen in Figure 1 is that WD 1121+216 is near a brighter source, and even a 2 pixel aperture on the white dwarf is contaminated by the light from the nearby star. Hence, we provide only an upper limit for $8 \mu\text{m}$ photometry of this object.

WD 2140+207 is a DQ star showing molecular C_2 Swan bands in the optical. The presence of carbon in cool He white dwarf atmospheres is thought to be the result of convective dredge-up (Pelletier et al. 1986). Figure 1 shows that the SED of WD 2140+207 fits a blackbody distribution reasonably well. The absence of metal lines in the optical spectrum of this star, plus the absence of excess infrared radiation, provide further evidence that the carbon was not accreted from the interstellar medium or a circumstellar debris disk. One caveat between our analysis and BLR01’s analysis is that using pure He model atmospheres, BLR01 estimated $T_{\text{eff}} = 8860 \pm 300$ K for this star. Fitting $VRJHK$ and 4.5 and $8 \mu\text{m}$ photometry of this star with a blackbody, we estimate the temperature of this star to be 9407 K. None of the other stars in our sample that are in common with the BLR01 sample show this discrepancy. Dufour et al. (2005) reanalyzed the DQ stars in the BLR01 sample and demonstrated that the inclusion of carbon in model atmosphere calculations reduces the estimated effective temperatures and surface gravities for DQ white dwarfs. They obtained an even lower T_{eff} measurement (8200 ± 250 K) for WD 2140+207.

4. DISCUSSION

Our *Spitzer* observations showed that all H-rich white dwarfs with $T_{\text{eff}} < 7000$ K show slight mid-infrared flux deficits. Having several stars with small deficits makes these deficits significant. Moreover, LHS 1126 shows significantly depressed mid-infrared fluxes relative to white dwarf models. J. H. Debes et al. (2005, private communication) have also found 10%–20% flux deficits ($\geq 3 \sigma$ significance) at 4.5 , 5.6 , and $8 \mu\text{m}$ in two DAZ white dwarfs with $T_{\text{eff}} = 6820$ and 7310 K. Combining the facts that WD 0552–041 (with a pure He atmosphere) does not show a clear flux deficit, and all H-rich white dwarfs cooler than 7000 K




FIG. 4.—SED of LHS 1126, along with a 5400 K blackbody (dotted line). White dwarf models with $T_{\text{eff}} = 5400$ K, $\log g = 7.9$, and $\log N(\text{He})/N(\text{H}) = 1.5$ (lower solid line) and $\log N(\text{He})/N(\text{H}) = 1$ (upper solid line) are also shown. The dashed line shows a power law with $\alpha = 2$. [See the electronic edition of the *Journal* for a color version of this figure.]

do exhibit mid-infrared flux deficits, LHS 1126 and WD 1748+708 are likely to have mixed H/He atmospheres. This is also consistent with Bergeron et al.’s (1994) and Wolff et al.’s (2002) analysis. However, one question remains to be answered: can these flux deficits be explained by CIA?

The atmospheres of M, L, and T dwarfs are also expected to exhibit CIA (Borysov et al. 1997). Roellig et al. (2004) obtained mid-infrared spectroscopy of M, L, and T dwarfs in the $5\text{--}15 \mu\text{m}$ range. Their Figure 2 shows that the observed spectra are in good agreement with the model atmosphere calculations, with only a few minor deviations. Hence, CIA calculations for low-density atmospheres seem to be accurate.

The near-infrared ($1\text{--}2 \mu\text{m}$) flux deficit in LHS 1126 was discovered earlier by Wickramasinghe et al. (1982). Bergeron et al. (1994) and BRL97 explained this deficit as CIA by molecular hydrogen due to collisions with helium and found the H/He ratio to be ~ 0.01 . Wolff et al. (2002) used Faint Object Spectrograph data plus optical and infrared photometry of LHS 1126 to model this star’s SED in the $0.2\text{--}2.2 \mu\text{m}$ range. They found that the hydrogen abundance reported by Bergeron et al. (1994) would result in an extremely strong $\text{Ly}\alpha$ absorption and that the SED is best fitted with an abundance ratio of $\text{H}/\text{He} = 3 \times 10^{-6}$. Wolff et al. (2002) and Bergeron et al. (1994) models do not give a consistent picture for the H/He ratio, and neither model is adequate. We revisit this problem by extending our wavelength coverage to $8 \mu\text{m}$. Figure 4 shows the ultraviolet spectrum (kindly made available to us by D. Koester) and optical and infrared photometry of LHS 1126 along with a 5400 K blackbody (dotted line). Mixed atmosphere white dwarf models (kindly made available to us by D. Saumon) with $T_{\text{eff}} = 5400$ K, $\log g = 7.9$, and $\log N(\text{He})/N(\text{H}) = 1.5$ (lower solid line) and $\log N(\text{He})/N(\text{H}) = 1$ (upper solid line) are also shown. These white dwarf models include CIA opacities but not $\text{Ly}\alpha$ absorption; therefore, they cannot be used to match the ultraviolet data. If we just use the optical and near-infrared photometry (as in Bergeron et al. 1994), we could fit the observations with a $\log N(\text{He})/N(\text{H}) = 1\text{--}1.5$ model. However, these models cannot explain the observed flux deficits in the *Spitzer* observations. Bergeron et al. (1995) and Hansen (1998) models cannot explain these flux deficits either (B. M. S. Hansen & P. Bergeron 2005, private communication). CIA is expected to create wiggles and

bumps in the spectrum up to $3 \mu\text{m}$, and the SEDs of cool white dwarfs are expected to return to normal in the mid-infrared (Fig. 4; see also Fig. 6 of Jorgensen et al. 2000).

Figure 4 also shows a power-law distribution with $\alpha = 2$, i.e., a Rayleigh-Jeans spectrum (*dashed line*). The infrared SED of LHS 1126 follows a Rayleigh-Jeans spectrum fairly well. It is striking to note that we can fit the ultraviolet and optical part of the SED with a 5400 K blackbody, and the infrared part of the SED with $T_{\text{eff}} > 10^5$ K. The power law is a reasonable, but not a perfect, fit to the infrared SED of this object, and photometry in the other IRAC channels or spectroscopy with *Spitzer* may reveal more structure in the mid-infrared, but the current data suggest the following:

1. The CIA opacity calculations are incomplete, and there is a significant, unexplained flux deficit in the mid-infrared. As the densities in white dwarf atmospheres approach a significant fraction of the liquid state densities, molecular absorption (H_2 - H_2 , H_2 -He), emission, and light scattering are expected to become increasingly important. Current CIA opacity calculations in dense media may be incomplete in certain wavelength regimes, for example, the rovibrational band of H_2 CIA may be enhanced compared to predictions (L. Frommhold & D. Saumon 2005, private communication). On the other hand, the featureless spectra of the ultracool white dwarfs (Gates et al. 2004), and the reasonably good fit of a Rayleigh-Jeans distribution to the LHS 1126 SED in the infrared, suggest another mechanism to explain the mid-infrared flux deficits observed in white dwarfs. CIA is expected to disappear with the dissociation of molecules when T_{eff} increases past 5500 K. Therefore, white dwarfs with $T_{\text{eff}} > 6000$ K that show infrared flux deficits could not be explained by changing the CIA opacities.

2. The mid-infrared flux deficits are caused by some as yet unrecognized physical process(es). The problem may be caused by an unknown or poorly understood absorption process, or it may be the result of a different source function operating in these dense atmospheres. H^- bound-free absorption is thought

to be the most important source of opacity in warmer DA white dwarfs. More work is required to test if possible changes in this absorption could help explain the mid-infrared flux deficits. The input physics used in white dwarf model atmospheres is mostly based on the ideal gas approximation. The extreme conditions in white dwarf atmospheres require a new look at dense medium effects on the equation of state, chemistry, opacities, and radiative transfer. Refractive opacities, presence of heavy elements, or formation of trace species and many other factors can change the opacity sources in white dwarf atmospheres (see Kowalski [2006] for a detailed discussion).

Near/mid-infrared photometry and spectroscopy of ultracool white dwarfs and the so-called C_2H stars will be useful to test these ideas. At this time, we only report and do not understand the observed mid-infrared flux deficits, but this mystery needs to be resolved before we can use the cool white dwarfs as accurate chronometers to find the age of Galactic populations.

We would like to thank Didier Saumon, Lothar Frommhold, Jason Kalirai, Brad Hansen, and Pierre Bergeron for helpful discussions, and our referee, Jay Farihi, for helpful suggestions that greatly improved the article. This work is based in part on observations made with the *Spitzer Space Telescope*, which is operated by the Jet Propulsion Laboratory (JPL), California Institute of Technology under NASA contract 1407. Support for this work was provided by NASA through award project NBR: 1269551 issued by JPL/California Institute of Technology to the University of Texas. This publication makes use of data products from the 2MASS, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by NASA and the NSF. This material is based upon work supported by NASA under grant NAG5-13070 issued through the Office of Space Science.

REFERENCES

- Adelman-McCarthy, J. K., et al. 2006, *ApJS*, 162, 38
 Angel, J. R. P. 1977, *ApJ*, 216, 1
 Becklin, E. E., et al. 2005, *ApJ*, 632, L119
 Bergeron, P., & Leggett, S. K. 2002, *ApJ*, 580, 1070
 Bergeron, P., Leggett, S. K., & Ruiz, M. T. 2001, *ApJS*, 133, 413 (BLR01)
 Bergeron, P., Ruiz, M. T., & Leggett, S. K. 1997, *ApJS*, 108, 339 (BRL97)
 Bergeron, P., et al. 1994, *ApJ*, 423, 456
 ———. 1995, *ApJ*, 443, 764
 Borysow, A., et al. 1997, *A&A*, 324, 185
 Dufour, P., Bergeron, P., & Fontaine, G. 2005, *ApJ*, 627, 404
 Farihi, J. 2004, *ApJ*, 610, 1013
 ———. 2005, *AJ*, 129, 2382
 Fazio, G. G., et al. 2004, *ApJS*, 154, 10
 Frommhold, L. 1993, *Collision-Induced Absorption in Gases* (Cambridge: Cambridge Univ. Press)
 Gates, E., et al. 2004, *ApJ*, 612, L129
 Hansen, B. M. S. 1998, *Nature*, 394, 860
 Hansen, B. M. S., et al. 2004, *ApJS*, 155, 551
 Harris, H. C., et al. 2001, *ApJ*, 549, L109
 ———. 2006, *AJ*, 131, 571
 Jorgensen, U. G., Hammer, D., Borysow, A., & Falcksgaard, J. 2000, *A&A*, 361, 283
 Kilic, M., von Hippel, T., Leggett, S. K., & Winget, D. E. 2005, *ApJ*, 632, L115
 Kilic, M., et al. 2006, *AJ*, 131, 582
 Kowalski, P. M. 2006, *ApJ*, in press
 Kowalski, P. M., & Saumon, D. 2004, *ApJ*, 607, 970
 Kowalski, P. M., Saumon, D., & Mazevet, S. 2005, in *ASP Conf. Ser.* 334, 14th European Workshop on White Dwarfs, ed. D. Koester & S. Moehler (San Francisco: ASP), 203
 Maxted, P. F. L., Marsh, T. R., & Moran, C. K. J. 2000, *MNRAS*, 319, 305
 McCook, G. P., & Sion, E. M. 1999, *ApJS*, 121, 1
 Monelli, M., et al. 2005, *ApJ*, 621, L117
 Oppenheimer, B. R., et al. 2001, *ApJ*, 550, 448
 Pelletier, C., et al. 1986, *ApJ*, 307, 242
 Reach, W. T., et al. 2005a, *ApJ*, 635, L161
 ———. 2005b, *PASP*, 117, 978
 Roellig, T. L., et al. 2004, *ApJS*, 154, 418
 Saumon, D., & Jacobson, S. B. 1999, *ApJ*, 511, L107
 Schmidt, G. D., Bergeron, P., & Fegley, B., Jr. 1995, *ApJ*, 443, 274
 Werner, M. W., et al. 2004, *ApJS*, 154, 1
 Wickramasinghe, D. T., Allen, D. A., & Bessel, M. S. 1982, *MNRAS*, 198, 473
 Wolff, B., Koester, D., & Liebert, J. 2002, *A&A*, 385, 995
 Zuckerman, B., & Becklin, E. E. 1987, *Nature*, 300, 138
 Zuckerman, B., Koester, D., Reid, I. N., & Hunsch, M. 2003, *ApJ*, 596, 477