

4-1979

Ca II H and K and Hydrogen-Line Variations in V471 Tauri (=BD + 16°516)

Terry D. Oswalt

Ohio State and Ohio Wesleyan Universities, oswalb32@erau.edu

Follow this and additional works at: <https://commons.erau.edu/publication>



Part of the [Stars, Interstellar Medium and the Galaxy Commons](#)

Scholarly Commons Citation

Oswalt, T. D. (1979). Ca II H and K and Hydrogen-Line Variations in V471 Tauri (=BD + 16°516). *Publications of the Astronomical Society of the Pacific*, 91(540). <https://doi.org/10.1086/130475>

This Article is brought to you for free and open access by Scholarly Commons. It has been accepted for inclusion in Publications by an authorized administrator of Scholarly Commons. For more information, please contact commons@erau.edu.

Ca II H AND K AND HYDROGEN-LINE VARIATIONS IN V471 TAURI (= BD +16°516)

TERRY D. OSWALT

Perkins Observatory, The Ohio State and Ohio Wesleyan Universities

Received 1978 December 11

Spectroscopic and narrow-band photoelectric observations of the white dwarf eclipsing binary V471 Tau (= BD +16°516) are reported. Periodic fluctuations in Ca II H and K emission and H α absorption, and a wave-like distortion of the light curve mimic the characteristics of several well-observed RS CVn binaries. A probable correlation between maximum emission-line strength and wave minimum is shown to exist.

Key words: eclipsing binaries—white dwarfs—emission-line variations

Introduction

The discovery of the eclipsing nature of BD +16°516 (= V471 Tauri) by Nelson and Young (1970) unleashed a brief flurry of excitement among observers and theorists. Not only was the opportunity for the determination of reliable physical data for a white dwarf anticipated, but early calculations indicated that the degenerate component was one of the hottest known (Hills 1971; Young and Nelson 1972). When ambiguities in the solution of the photometric light curve due to the very small value for the ratio of the radii (~ 0.01) became apparent, interest waned. Within the last few years, however, this system has again received attention because of the cooler component's striking similarities to many of the so-called RS Canum Venaticorum variables, which exhibit peculiar migrating wave-like light-curve distortions but do not contain degenerate components. V471 Tau is known to exhibit such a photometric distortion wave, migrating toward decreasing orbital phase with a period of about one year.

Nelson and Young (1976) have published a concise review of work that has been done to date on this system. Hall (1976) has presented a comprehensive discussion of the RS CVn variables. Some of the similarities of V471 Tau to this class of binaries were specifically mentioned.

Photoelectric *UBV*, *woby*, H α , and H β photometry, and spectroscopic work conducted during the last three observing seasons at the Perkins and Lowell Observatories has revealed additional similarities between V471 Tau and several of the most active RS CVn variables (Oswalt 1978). Presented here are the results of the narrow-band photometry and spectroscopic work. Concurrent *UBV* and *woby* light curves obtained on all nights will be published elsewhere.

Equipment and Observations

The spectrograms were obtained at the Perkins 1.8-

m telescope of the Ohio State and Ohio Wesleyan Universities in Flagstaff, Arizona, equipped with the Boller and Chivens single-stage image-tube spectrograph. All the exposures were made on unbaked IIA-D emulsion at a reciprocal dispersion of about 77 \AA mm^{-1} near the 4640 \AA blaze wavelength of the grating.

The H β photometry was obtained with the single-channel OSU "red" photometer attached to the 0.8-m telescope at Perkins Observatory in Delaware, Ohio. A refrigerated EMI 6256 S-11 photomultiplier operated in the pulse-counting mode was used. The H β filter pair used closely matches the transmission characteristics defined by Crawford and Mander (1966).

A similar photometer was used for the H α observations. These were obtained with the 1.1-m and Perkins 1.8-m telescopes at Lowell Observatory in Flagstaff. Refrigerated EMI 9558 S-20 and ITT F4085 S-20 photomultipliers operated in the pulse-counting mode were used for the 1976 and 1977 observations, respectively. Bandpass half-widths for the narrow and wide filters are about 28 \AA and 150 \AA , respectively.

The primary comparison star was BD +17°638 (= HD 24040). A check star, BD +16°544, was monitored occasionally on several nights. No variations in either of these stars were noted.

An observing sequence SCVCS was followed throughout each night. On those nights when the check star was observed readings were taken at least once an hour. Integration times were integral multiples of ten seconds. Three or more such readings were averaged for an observation, depending on the count rates in a particular filter. The standard error of a single observation, as determined from the internal scatter of the comparison and variable star readings, was considerably better than ~ 0.015 in all bandpasses for most nights. All data have been corrected for atmospheric extinction. Phases of all observations are calculated from the heliocentric linear ephemeris given by Young and Lanning (1975).

Spectroscopic Results

During a separate observing program 17 image-tube spectrograms were obtained between 1975 October 6–11. Medium-density exposures ranged from three to seven minutes depending on sky conditions and were grouped mostly about phase 0.5 and eclipse for the purpose of detecting any transient spectral features during ingress and egress. These plates were examined in detail for spectral variations. At least during this time the emission components of Ca II H and K were strongly variable, *apparently* in synchronism with the orbital period. Maximum emission levels occurred near phase 0.5, with virtually zero emission during eclipse totality. Density tracings from the most uniformly exposed plates are shown in Figure 1.

More complete phase coverage was obtained during the period 1976 January 1–6. These 14 spectra covered about 80% of the 12.5-hour orbital period. Ten exposures taken on January 5 provided continuous coverage between phase ~ 0.5 and eclipse. The same periodic variability in Ca II H and K emission was evident: maximum near phase 0.5, minimum near eclipse. Two plates in particular taken during ingress showed a distinct strong double-emission core in the K line. No other obvious emission features are present in this collection of spectra, except O I $\lambda 5577$ due to the night sky.

Narrow-Band Photometry

A search for absorption-line changes was made and H β seemed to be marginally variable (Fig. 2). Consequently, H β photometry was undertaken during the following season with the 0.8-m telescope at Perkins Observatory. These observations are presented in Table I. The instrumental H β index is defined conventionally:

$$\beta' = -2.5 \log F(\text{H}\beta)_n / F(\text{H}\beta)_w,$$

where $F(\text{H}\beta)_n$ and $F(\text{H}\beta)_w$ are the observed fluxes through the narrow and wide H β filters, respectively. The concurrent V and β' observations are presented in Figure 3. The migrating nature of the wave is clearly evident in such composite V light curves, however no periodic variability above the noise greater than about 0^m02 in β' was detected.

It was presumed that any real periodic variations in H β would be accompanied by larger fluctuations in H α . Later in the same season H α photometry was begun at the Lowell 1.1-m and Perkins 1.8-m telescopes. These data are listed in Table II, where an instrumental H α index is defined in a manner similar to β' :

$$\alpha' = -2.5 \log F(\text{H}\alpha)_n / F(\text{H}\alpha)_w.$$

Figure 4a presents the V light curve for these four nights. It is virtually identical in appearance to those

obtained in the wide bandpass H α filter. Immediately below this is a plot of the behavior of α' with phase for comparison (Fig. 4b). The 1976 observations indicate $\sim 0^m06$ variation in α' ; a "dip" in H α flux (increase in α') occurs at about phase 0.0 in Figure 4b. This may be interpreted to arise from the same mechanism producing the fluctuations in Ca II H and K emission. It can be supposed that a small emission component partially filled in the H α absorption line outside eclipse. Short-exposure, high-dispersion line profiles would be required to confirm this. Interestingly, Lanning and Etzel (1976) have reported the detection of H α emission in V471 Tau on at least one occasion.

It was not apparent from the 1976 observations that the variations in H α had anything at all to do with the photometric distortion wave, since wave maximum occurred about 1/4 cycle before the "dip" in H α flux (maximum α'), and wave minimum about 0.6 earlier in phase than the "dip". Besides, minimum H and K emission and H α flux appeared to occur during eclipse totality. Rather, some sort of excitation due to the hot companion outside eclipse seemed probable. Another interesting feature of the 1976 V light curve was that observations made during wave minimum seemed to exhibit considerably more scatter than elsewhere in the light curve, even though the individual ten-second readings averaged to obtain each observation were consistent to better than 1%.

Additional H α observations were obtained at Lowell Observatory during December 1977. These are presented along with the 1976 data in Table II, plotted versus phase with the 1976 data in Figure 4c, and plotted separately in Figure 4d. The first new nights in 1977 were of generally lower photometric quality than the previous year, however α' values for the comparison and check stars were found to be virtually independent of the small changes in sky transparency which affected the long-term stability of individual filter readings. This is evident by the comparable intrinsic scatter in Figures 4b and 4d.

The 1977 observations indicate that either the "dip" in H α flux was much broader than was apparent during the previous year, or that it had migrated toward decreasing orbital phase. Observations scheduled during the 1978 season should help decide between these alternatives. If the latter possibility is correct this shift is in the same sense as that observed for the distortion wave in the broad-band light curves. In addition, the "dip" in H α flux still would have occurred about 1/4 cycle past wave maximum. It is not obvious why such a phase lag might occur, but some sort of aspect effect is suggested. The incomplete phase coverage of the 1977 data makes it difficult to judge whether the mean




FIG. 1—Density tracings of spectrograms showing the behavior of Ca II H and K emission with orbital phase in V471 Tau during October 1975. Numbers along right side are UT day of exposures. Density scale is arbitrary but similar for all tracings.

systematic α' changed between the two seasons although the “dip” appears to be $\sim 0^m.03$ shallower than in 1976.

Minimum $H\alpha$ flux (maximum α') seems to occur at or near the base of the descending branch of wave

maximum, immediately before the most level portion of the light curve. The peak of the broad asymmetrical $H\alpha$ flux maximum (α' minimum) evidently corresponded with wave minimum in November 1976.

The incomplete 1977 observations seem to indicate




FIG. 2—H β absorption-line variations in V471 Tau during October 1975. Emission lines near yellow end of spectrum (right) are due to night sky.




FIG. 3—Photometric behavior of V471 Tau in 1976: Differential V magnitudes (top), and β' observations (bottom). Standard error of a single observation is comparable to symbol size.

that the phase of maximum H α emission does not always coincide with wave minimum, however. This suggests that rather long-lived emitting regions may migrate in longitude with respect to the darkened areas presumed responsible for the distortion wave minimum. Such behavior is that which might be expected from large-scale prominence activity arising from the active regions, particularly if differential rotation is taking place as supposed in the RS CVn variables (Hall 1972). There exists strong circumstantial evidence for such chromospheric events in several of these binaries (Weiler 1978*b*) and in V471 Tau (Young and Lanning 1975; Young 1976). This interpretation may shed some light on the conflicting phase correlations between emission-line strengths and the distortion wave observed in some

of these stars (Droppo and Milone 1976). At any rate, it appears that the variations in H α are related to the distortion wave in the same sense that Weiler (1978*a*) found for Ca II H and K emission in several RS CVn variables, and not exclusively the result of excitation by the hot companion.

The Light Curve Variations

No unique model has yet been presented which explains the peculiarities of the light-curve distortion wave exhibited by V471 Tau, although like the RS CVn binaries, several characteristics of the system can be explained very simply by relatively dark regions rotating with the surface of the cooler component: variability of eclipse depths, changes in wave amplitude,

TABLE I

H β Observations of V471 Tauri		
Julian Date (2,440,000+)	Phase	β'
3053.665	0.560	1.323
3053.686	0.601	1.349
3053.721	0.668	1.388
3053.743	0.710	1.363
3053.763	0.749	1.392
3053.784	0.789	1.380
3053.807	0.833	1.393
3053.831	0.878	1.384
3053.850	0.916	1.412
3053.869	0.952	1.397
3053.885	0.982	1.394
3053.903	0.018	1.386
3083.755	0.294	1.348
3083.772	0.328	1.332
3083.796	0.373	1.374
3083.814	0.407	1.378
3083.835	0.448	1.386
3083.847	0.472	1.371
3083.873	0.522	1.345
3083.897	0.566	1.383
3083.910	0.591	1.377
3083.931	0.632	1.356
3083.944	0.656	1.376
3099.718	0.924	1.404
3099.735	0.955	1.378
3099.759	0.002	1.368
3099.780	0.041	1.391
3099.799	0.078	1.405
3099.811	0.101	1.365
3099.830	0.138	1.380
3099.841	0.159	1.370
3099.866	0.207	1.411
3099.879	0.231	1.382
3099.902	0.276	1.342
3102.670	0.586	1.396
3102.687	0.620	1.379
3102.704	0.652	1.375
3102.721	0.684	1.384
3102.733	0.707	1.390
3102.754	0.747	1.394
3102.821	0.877	1.352
3103.545	0.265	1.382
3103.561	0.297	1.375
3103.581	0.335	1.352
3103.597	0.366	1.379
3127.621	0.460	1.350
3127.637	0.491	1.347
3127.649	0.514	1.342
3127.663	0.542	1.371
3127.678	0.571	1.364

TABLE I (Continued)

Julian Date (2,440,000+)	Phase	β'
3127.691	0.594	1.388
3127.707	0.626	1.379
3127.724	0.658	1.406
3127.737	0.683	1.395
3127.775	0.757	1.381

migration of the wave, and, to a certain extent, apparent changes in period. That the wave-like variations are intrinsic to the cooler component is quite certain: the 1977 light curves show clear continuation of the wave during eclipse of the hot companion.

Spotted star models such as those developed by Friedemann and Gürtler (1975) and Vogt (1975) predict a quasi-sinusoidal distortion wave if the darkened region extends over 180° in longitude, and flat maxima (minima) if less (more) than one hemisphere of the star is darkened. Some of the light curves of V471 Tau obtained by Ibanoglu (1976) and possibly Cester and Pucillo (1976) may be explained by the first situation. Frequently, however, V471 Tau exhibits a temporary constant light level between wave extrema. Moreover, this "reference" or "R-level" does not always bisect the wave; wave maximum often has a larger amplitude relative to this level than does wave minimum. Occasionally the situation is reversed. In addition, V471 Tau at times exhibits little evidence of a distortion wave at all (Oswalt 1978).

The published *V* light curves of V471 Tau which exhibit a well-defined R-level between wave maximum and minimum seem to share a common characteristic: the magnitude and color of this R-level seems to be fairly constant from season to season despite striking changes in the distortion wave on time-scales as short as several weeks. This is especially evident among those observations made over several seasons on the same or similar instrumental systems (Young and Nelson 1972; Ibanoglu 1976; Oswalt 1978). Such behavior suggests the occasional existence of a relatively spot-free hemisphere on the cool component. In this case the R-level would provide the most appropriate basis for determination of the photometric properties of the system, particularly when it coincides with the eclipse of the degenerate component.

It is interesting, though not unprecedented, that the above interpretation of the R-level would require co-existent bright and dark regions to explain the distortion wave maximum and minimum, respectively. Similar phenomena were suggested as an explanation for the light curve irregularities observed in AR Lacertae and YY Geminorum (Kron 1947, 1952). In addition, transient hot spots have been strongly indicated for the

TABLE II

TABLE II (Continued)

H α Observations of V471 Tauri 1976			Julian Date (2,440,000+)	Phase	α'
Julian Date (2,440,000+)	Phase	α'			
3107.728	0.291	1.486	3110.869	0.319	1.479
3107.743	0.320	1.457	3110.876	0.331	1.456
3107.756	0.346	1.462	3110.892	0.362	1.468
3107.769	0.371	1.459	3110.902	0.381	1.474
3107.786	0.403	1.487	3110.920	0.416	1.441
3107.810	0.449	1.474	3110.941	0.456	1.455
3107.825	0.479	1.471	3110.953	0.480	1.466
3107.843	0.512	1.467	3110.969	0.509	1.455
3107.854	0.534	1.460	3110.982	0.534	1.452
3107.869	0.561	1.488	3111.001	0.571	1.486
3107.885	0.592	1.464	3111.686	0.886	1.487
3107.898	0.618	1.488	3111.698	0.908	1.478
3107.914	0.648	1.476	3111.704	0.919	1.514
3107.940	0.698	1.488	3111.712	0.936	1.489
3107.956	0.729	1.493	3111.792	0.089	1.518
3108.676	0.111	1.518	3111.801	0.106	1.510
3108.687	0.132	1.488	3111.810	0.124	1.465
3108.697	0.151	1.492	3111.823	0.149	1.484
3108.709	0.174	1.501	3111.832	0.166	1.506
3108.719	0.193	1.483	3111.844	0.190	1.496
3108.730	0.213	1.481	3111.853	0.206	1.486
3108.734	0.222	1.475	3111.865	0.230	1.521
3108.818	0.382	1.477	3111.874	0.247	1.490
3108.827	0.401	1.488	3111.888	0.273	1.493
3108.834	0.414	1.487	3111.894	0.285	1.459
3108.842	0.429	1.478	3111.916	0.327	1.456
3108.857	0.458	1.474	3111.932	0.358	1.459
3108.867	0.477	1.480	3111.945	0.382	1.476
3108.880	0.501	1.470	3111.954	0.400	1.445
3108.889	0.520	1.467	3111.967	0.426	1.448
3108.899	0.539	1.477	3111.987	0.463	1.511
3108.911	0.560	1.481	3111.999	0.486	1.474
3108.922	0.583	1.472	3112.010	0.508	1.470
3108.932	0.601	1.487	3112.022	0.531	1.489
3108.943	0.624	1.479			
3108.950	0.636	1.508	3481.748	0.928	1.551
3108.961	0.658	1.478	3481.768	0.965	1.506
3110.667	0.930	1.506	3481.778	0.984	1.506
3110.678	0.952	1.513	3481.812	0.050	1.542
3110.692	0.979	1.514	3481.882	0.185	1.512
3110.702	0.999	1.526	3482.675	0.705	1.481
3110.705	0.004	1.526	3482.686	0.726	1.497
3110.714	0.020	1.517	3482.700	0.755	1.481
3110.716	0.025	1.468	3482.721	0.795	1.525
3110.726	0.043	1.530	3482.752	0.854	1.555
3110.786	0.160	1.475	3482.775	0.898	1.546
3110.804	0.193	1.461	3482.824	0.992	1.527
3110.823	0.230	1.450	3482.879	0.097	1.510
3110.836	0.254	1.432	3482.908	0.153	1.516
3110.847	0.277	1.489	3482.929	0.193	1.522

TABLE II (Continued)

Julian Date (2,440,000+)	1977 Phase	α'
3482.958	0.249	1.506
3483.624	0.527	1.505
3483.634	0.546	1.498
3483.649	0.575	1.507
3483.658	0.593	1.488
3483.669	0.613	1.484
3483.683	0.641	1.503
3483.694	0.662	1.508
3483.708	0.689	1.511
3483.722	0.716	1.514
3483.731	0.732	1.514
3483.744	0.758	1.543
3483.772	0.810	1.539
3483.806	0.876	1.560
3483.815	0.894	1.546
3483.826	0.915	1.514
3483.836	0.934	1.564
3483.851	0.963	1.560
3483.861	0.982	1.536
3483.866	0.992	1.542
3483.878	0.015	1.532
3483.883	0.024	1.542
3483.894	0.044	1.530
3483.906	0.069	1.547
3483.917	0.090	1.541
3483.927	0.109	1.515
3483.936	0.125	1.546
3483.961	0.174	1.481

flare star BY Draconis (Fix and Spangler 1976).

The absence of strong Balmer emission and flare activity would seem to imply that the cool component of V471 Tau displays a somewhat less-active chromo-

spheric network than either the most active RS CVn variables or flare stars. Continuing observations are planned for V471 Tau, and hopefully a more complete model of this interesting system can eventually be constructed.

It is a pleasure to thank Dr. Nathaniel White who generously lent his time and photoelectric equipment during the 1976 and 1977 observing runs at Lowell Observatory. Dr. Terry Roark kindly obtained the image-tube spectra concurrently with the author's 1976 photometry. Several helpful and informative discussions with Dr. Gerald Newsom and comments by an anonymous referee are also gratefully acknowledged.

REFERENCES

- Cester, B., and Pucillo, M. 1976, *Astr. and Ap.* 46, 197.
 Crawford, D. L., and Mander, J. 1966, *A.J.* 71, 114.
 Droppo, J. L., and Milone, E. F. 1976, *Inf. Bull. Var. Stars* No. 1130.
 Fix, J. D., and Spangler, S. R. 1976, *Ap. J. (Letters)* 205, L163.
 Friedemann, C., and Gürtler, J. 1975, *A.N. Bd.* 296, 125.
 Hall, D. S. 1972, *Pub. A.S.P.* 84, 323.
 ——— 1976, in *Multiple Periodic Variable Stars*, I.A.U. Colloquium No. 29, W. S. Fitch, ed. (Dordrecht: D. Reidel Publ. Co.), p. 287.
 Hills, J. G. 1971, *Pub. A.S.P.* 83, 81.
 Ibanoglu, C. 1976, *Inf. Bull. Var. Stars* No. 1088.
 Kron, G. E. 1947, *Pub. A.S.P.* 59, 261.
 ——— 1952, *Ap. J.* 115, 301.
 Lanning, H. H., and Etzel, P. B. 1976, *Inf. Bull. Var. Stars* No. 1147.
 Nelson, B., and Young, A. 1970, *Pub. A.S.P.* 82, 699.
 ——— 1976, in *Structure and Evolution of Close Binary Stars*, I.A.U. Symposium No. 73, P. Eggleton, S. Mitton and J. Whelan, eds. (Dordrecht: D. Reidel Publishing Co.), p. 141.
 Oswalt, T. D. 1978, *Bull. A.A.S.* 10, 410.
 Vogt, S. S. 1975, *Ap. J.* 199, 418.
 Weiler, E. J. 1978a, *M.N.R.A.S.* 182, 77.
 ——— 1978b, *A.J.* 83, 795.
 Young, A. 1976, *Ap. J.* 205, 182.
 Young, A., and Lanning, H. H. 1975, *Pub. A.S.P.* 87, 461.
 Young, A., and Nelson, B. 1972, *Ap. J.* 173, 653.




FIG. 4—Photometric observations of V471 Tau: (a) 1976 V light curve, (b) 1976 α' variations, (c) composite 1976-77 α' plot, (d) 1977 α' variations, (e) 1977 V light curve. Symbol size is comparable to standard error of a single observation. In all plots open symbols represent 1976 observations, solid symbols those made in 1977.