

3-2007

The Complex Cubi-Quintic Ginzburg-Landau Equation: Hopf Bifurcations Yielding Traveling Waves

Stefan C. Mnacas

University of Central Florida, mancass@erau.edu

S. Roy Choudhury

University of Central Florida

Follow this and additional works at: <https://commons.erau.edu/publication>



Part of the [Applied Mathematics Commons](#)

Scholarly Commons Citation

Mnacas, S. C., & Choudhury, S. R. (2007). The Complex Cubi-Quintic Ginzburg-Landau Equation: Hopf Bifurcations Yielding Traveling Waves. *Mathematics and Computers in Simulation*, 74(4/5).
<https://doi.org/10.1016/j.matcom.2006.10.022>

This Article is brought to you for free and open access by Scholarly Commons. It has been accepted for inclusion in Publications by an authorized administrator of Scholarly Commons. For more information, please contact commons@erau.edu.

The Complex Cubic–Quintic Ginzburg–Landau Equation: Hopf Bifurcations Yielding Traveling Waves

Stefan C. Mancas and S. Roy Choudhury

Department of Mathematics

University of Central Florida

Orlando, FL 32816–1364

mancass@erau.edu

choudhur@longwood.cs.ucf.edu

Abstract

In this paper we use a traveling wave reduction or a so-called spatial approximation to comprehensively investigate the periodic solutions of the complex cubic–quintic Ginzburg–Landau equation. The primary tools used here are Hopf bifurcation theory and perturbation theory. Explicit results are obtained for the post–bifurcation periodic orbits and their stability. Generalized and degenerate Hopf bifurcations are also briefly considered to track the emergence of global structure such as homoclinic orbits.

1 Introduction

The cubic complex Ginzburg–Landau equation (CGLE) is the canonical equation governing the weakly nonlinear behavior of dissipative systems in a wide variety of disciplines [11]. In fluid mechanics, it is also often referred to as the Newell–Whitehead equation after the authors who derived it in the context of Bénard convection [11, 17].

As such, it is also one of the most widely studied nonlinear equations. Many basic properties of the equation and its solutions are reviewed in [4, 8], together with applications to a vast variety of phenomena including nonlinear waves, second–order phase transitions, superconductivity, superfluidity, Bose–Einstein condensation, liquid crystals and string theory. The numerical studies by Brusch et al. [10, 9] which primarily consider periodic traveling wave solutions of the cubic CGLE, together with secondary pitchfork bifurcations and period doubling cascades into disordered turbulent regimes, also give comprehensive summaries of other work on this system. Early numerical studies [22, 23] and theoretical investigations [26, 27] of periodic solutions and secondary bifurcations are also of general interest for our work here.

Certain situations or phenomena, such as where the cubic nonlinear term is close to zero, may require the inclusion of higher–order nonlinearities leading to the so-called cubic–quintic CGLE [34]. This has proved to be a rich system with very diverse solution behaviors. In

particular, a relatively early and influential review by van Saarloos and Hohenberg, also recently extended to two coupled cubic CGL equations [33, 3], considered phase–plane counting arguments for traveling wave coherent structures, some analytic and perturbative solutions, limited comparisons to numerics, and so–called “linear marginal stability analysis” to select the phase speed of the traveling waves.

Among the multitude of other papers, we shall only refer to two sets of studies which will directly pertain to the work in this article. The first class of papers [21, 15, 14] and [18, 13] used dynamical systems techniques to prove that the cubic–quintic CGLE admits periodic and quasi–periodic traveling wave solutions.

The second class of papers [32, 2], primarily involving numerical simulations of the full cubic–quintic CGL PDE in the context of Nonlinear Optics, revealed various branches of plane wave solutions which are referred to as continuous wave (CW) solutions in the Optics literature. More importantly, these latter studies also found various spatially confined coherent structures of the PDE, with envelopes which exhibit complicated temporal dynamics. In [2], these various structures are categorized as plain pulses (or regular stationary solutions), pulsating solitary waves, creeping solitons, slugs or snakes, erupting solitons, and chaotic solitons depending on the temporal behavior of the envelopes. In addition, note that the speed of the new classes of solutions may be zero, constant, or periodic (since it is determined by boundary conditions, the speed is an eigenvalue, and it may be in principle also quasiperiodic or chaotic, although no such cases appear to have been reported). All indications are that these classes of solutions, all of which have amplitudes which vary in time, do not exist as stable structures in Hamiltonian systems. Even if excited initially, amplitude modulated solitary waves restructure into regular stationary solutions [6]. Exceptions to this rule are the integrable models where the pulsating structures are nonlinear superpositions or fundamental solutions [31]. Hence, these classes of solutions are novel and they exist only in the presence of dissipation in the simulations of [2]. Also, secondary complete period doubling cascades of the pulsating solitons leading as usual to regimes of chaos are also found. This last feature for numerical solutions of the full cubic–quintic PDE is strongly reminiscent of the period doubling cascades found in [10, 9] for period solutions of the traveling wave reduced ODEs of the cubic CGLE.

Motivated by the above, we begin a fresh look at the traveling wave solutions of the cubic–quintic CGLE in this paper. Besides attempting to understand the complex numerical coherent structures in [2], one other goal is to build a bridge between the dynamical systems approach in [21, 15, 14]–[18, 13] and the numerical one in [32, 2]. Given the importance of the cubic–quintic CGLE as a canonical pattern–forming system, this is clearly important in and of itself. However, a word of warning is in order here. Some of the features in [2] may well be inherently spatio–temporal in nature, so that a spatial traveling–wave reduction may not be sufficient to completely capture all aspects. Indeed, there is some evidence along these lines [6].

In this paper we begin by using a traveling wave reduction or a so–called spatial approximation to investigate the periodic solutions of the CGLE. The primary tool used here is Hopf bifurcation theory. Immediately following the Hopf bifurcations we construct the periodic orbits by using the method of multiple scales.

The remainder of the paper is organized as follows. We first analyze the stability of fixed points in Section 2 and the onset of instability via a Hopf bifurcation, which may be either supercritical or subcritical. Then stability of periodic orbits is presented in Section 3 where

we derive analytical expressions for the periodic orbits resulting from this Hopf bifurcation, and for their stability coefficients, by employing the multiple scales method. Section 4 considers numerical solutions and discusses the results. Generalized and degenerate Hopf bifurcations have also been considered to track the emergence of global structure such as homoclinic orbits in Section 5.

2 Stability Analysis of Fixed Points

In this section, we conduct a stability analysis of individual plane wave solutions using regular phase plane techniques. This was already done for the alternative formulation of the traveling wave ODEs given in [34]. We provide a brief derivation for our system (2.4a), (2.4b) and (2.4c) for completeness and future use. However, a much more complex question is the issue of categorizing and elucidating the possible existence of, and transitions among, multiple plane wave states which may co-exist for the same parameter values in (2.1) (corresponding to the same operating conditions of the underlying system). Such behavior is well-documented in systems such as the Continuous Stirred Tank Reactor System [31, 1]. For a system such as (2.1) and the associated ODEs (2.4a), (2.4b) and (2.4c), the large number of parameters makes a comprehensive parametric study of co-existing states bewilderingly complex, if not actually impracticable.

We shall consider the cubic-quintic CGLE in the form [34]

$$\partial_t A = \epsilon A + (b_1 + ic_1)\partial_x^2 A - (b_3 - ic_3)|A|^2 A - (b_5 - ic_5)|A|^4 A \quad (2.1)$$

noting that any three of the coefficients (no two of which are in the same term) may be set to unity by appropriate scalings of time, space and A .

For the most part, we shall employ the polar form used in earlier treatments [10, 34] of the traveling wave solutions of (2.1). This takes the form of the ansatz

$$\begin{aligned} A(x, t) &= e^{-i\omega t} \hat{A}(x - vt) \\ &= e^{-i\omega t} a(z) e^{i\phi(z)} \end{aligned} \quad (2.2)$$

where ω , a , and ϕ are all real,

$$z \equiv x - vt \quad (2.3)$$

is the traveling wave variable and ω and v are the frequency and translation speed (and are determined by the boundary conditions, and are thus eigenvalues of the uniformly translated solutions). Substitution of (2.2)/(2.3) in (2.1) leads, after some simplification, to the three mode dynamical system

$$a_z = b \quad (2.4a)$$

$$b_z = a\psi^2 - \gamma_1 \left[\gamma_2 a + v \left(b_1 b + c_1 \psi a \right) - \gamma_3 a^3 - \gamma_4 a^5 \right] \quad (2.4b)$$

$$\psi_z = -\frac{2\psi b}{a} + \gamma_1 \left[\gamma_5 + v \left(\frac{c_1 b}{a} - b_1 \psi \right) - \gamma_6 a^2 - \gamma_7 a^4 \right] \quad (2.4c)$$

where $\psi \equiv \phi_z$. Note that we have put the equations into a form closer to that in [10], rather than that in [34], so that (2.4) is a generalization of the traveling wave ODEs in [10] to

include the quintic terms and the constant terms $\gamma_1 - \gamma_7$ are given as functions of the system parameters in the following manner:

$$\begin{aligned}\gamma_1 &= \frac{1}{b_1^2 + c_1^2} \\ \gamma_2 &= b_1\epsilon + c_1\omega \\ \gamma_3 &= b_1b_3 - c_1c_3 \\ \gamma_4 &= b_1b_5 - c_1c_5 \\ \gamma_5 &= -b_1\omega + c_1\epsilon \\ \gamma_6 &= b_1c_3 + c_1b_3 \\ \gamma_7 &= b_1c_5 + c_1b_5 .\end{aligned}$$

From (2.2), a fixed point $(a_0, 0, \psi_0)$ of (2.4) corresponds to a plane wave solution

$$A(x, t) = a_0 e^{i(\psi_0 z - \omega t) + i\theta} \quad (2.5)$$

with θ an arbitrary constant.

The fixed points of (2.4a), (2.4b) and (2.4c) may be obtained by setting $b = 0$ (from (2.4a)) in the right hand sides of the last two equations, solving the last one for ψ , and substituting this in the second yielding the quartic equation

$$\alpha_4 x^4 + \alpha_3 x^3 + \alpha_2 x^2 + \alpha_1 x + \alpha_0 = 0 \quad (2.6)$$

with

$$x = a^2, \quad (2.7a)$$

$$\alpha_4 = \frac{\gamma_7^2}{b_1^2 v^2} \quad (2.7b)$$

$$\alpha_3 = \frac{2\gamma_6\gamma_7}{b_1^2 v^2} \quad (2.7c)$$

$$\alpha_2 = \frac{\gamma_6^2 - 2\gamma_5\gamma_7}{b_1^2 v^2} + \frac{\gamma_1(b_1\gamma_4 + c_1\gamma_7)}{b_1} \quad (2.7d)$$

$$\alpha_1 = \gamma_1\left(\gamma_3 + \frac{c_1\gamma_6}{b_1}\right) - \frac{2\gamma_5\gamma_6}{b_1^2 v^2} \quad (2.7e)$$

$$\alpha_0 = \frac{\gamma_5^2}{b_1^2 v^2} - \frac{\gamma_1}{b_1}(b_1\gamma_2 + c_1\gamma_5) . \quad (2.7f)$$

Thus, with $a_0 = \sqrt{x}$ for x any of the four roots of (2.6), we have a plane wave solution of the form (2.5). For each of the four roots $x_i, i = 1, \dots, 4$ of (2.6) corresponding to a fixed point of (2.4a), (2.4b) and (2.4c) or a plane wave $\sqrt{x_i} e^{i(\psi_i z - \omega t) + i\theta_i}$, the stability may be determined using regular phase-plane analysis. The characteristic polynomial of the Jacobian matrix of a fixed point $x_i = a_i^2$ of (2.4a), (2.4b) and (2.4c) may be expressed as

$$\lambda^3 + \delta_1 \lambda^2 + \delta_2 \lambda + \delta_3 = 0 \quad (2.8)$$

where

$$\delta_1 = 2b_1v\gamma_1 \tag{2.9a}$$

$$\delta_2 = 3\psi^2 + \gamma_1[\gamma_2 - a^2(3\gamma_3 + 5a^2\gamma_4) - v(3c_1\psi - v)] \tag{2.9b}$$

$$\begin{aligned} \delta_3 = & -2a^2\gamma_1(\gamma_6 + 2a^2\gamma_7)(-2\psi + c_1\gamma_1v) \\ & + b_1\gamma_1v[-\psi^2 + \gamma_1(\gamma_2 - 3a^2\gamma_3 - 5a^4\gamma_4 + c_1\psi v)] \end{aligned} \tag{2.9c}$$

where the fixed point values $(a_i, \psi_i) = (\sqrt{x_i}, \psi_i)$ are to be substituted in terms of the system parameters. Note that ψ_i is obtained by setting $a = a_i = \sqrt{x_i}$, and $b = 0$ in the right side of (2.4c).

For $(a_0, 0, \psi_0)$ to be a stable fixed point within the linearized analysis, all the eigenvalues must have negative real parts. Using the Routh–Hurwitz criterion, the necessary and sufficient conditions for (2.8) to have $Re(\lambda_{1,2,3}) < 0$ are:

$$\delta_1 > 0, \quad \delta_3 > 0, \quad \delta_1\delta_2 - \delta_3 > 0. \tag{2.10}$$

Equation (2.10) is thus the condition for stability of the plane wave corresponding to x_i .

On the contrary, one may have the onset of instability of the plane wave solution occurring in one of two ways. In the first, one root of (2.8) (or one eigenvalue of the Jacobian) becomes non-hyperbolic by going through zero for

$$\delta_3 = 0. \tag{2.11}$$

Equation (2.11) is thus the condition for the onset of “static” instability of the plane wave. Whether this bifurcation is a pitchfork or transcritical one, and its subcritical or supercritical nature, may be readily determined by deriving an appropriate canonical system in the vicinity of (2.11) using any of a variety of normal form or perturbation methods [16, 25, 7].

One may also have the onset of dynamic instability (“flutter” in the language of Applied Mechanics) when a pair of eigenvalues of the Jacobian become purely imaginary. The consequent Hopf bifurcation at

$$\delta_1\delta_2 - \delta_3 = 0 \tag{2.12}$$

leads to the onset of periodic solutions of (2.4a), (2.4b) and (2.4c) (dynamic instability or “flutter”). Note that the case $\delta_1 = 0$ is not enough to produce instability. These periodic solutions for $a(z)$ and $\psi(z)$, which may be stable or unstable depending on the super- or subcritical nature of the bifurcation, correspond via (2.2) to solutions

$$A(x, t) = a(z)e^{i(\int \psi dz - \omega t)} \tag{2.13}$$

of the CGLE (2.1) which are, in general, quasiperiodic wavetrain solutions. This is because the period of ψ and ω are typically incommensurate. Eq. (2.13) is periodic if $\omega = 0$.

3 Stability Analysis of Periodic Orbits

In this section we will use the method of multiple scales to construct analytical approximations for the periodic orbits arising through Hopf bifurcation of the fixed point of the CGLE equation. For the systems of differential equations given by (2.4a), (2.4b) and (2.4c), the

physically relevant point is given by $(a_0, 0, \psi_0)$ where ψ_i is obtained by setting $a = a_i = \sqrt{x_i}$, in

$$\psi_i = \frac{\gamma_5 - a_i^2(\gamma_6 + a_i^2\gamma_7)}{b_1 v} \quad (3.1)$$

and x_i is one of the roots of the fixed point equation (2.6). We will choose the parameter ϵ which represents the linear gain or loss as the control parameter. The limit cycle is determined by expanding about the fixed point using progressively slower spatial scales. In the standard way, we write the various or multiple scales as $z = Z_0, Z_1 = \delta Z_0, Z_2 = \delta^2 Z_0, \dots$, where δ is the usual multiple scales expansion parameter. We shall expand in powers of δ , to separate the various scales, and then set $\delta = 1$ at the end in the usual way. The expansion takes the form

$$a = a_0 + \sum_{n=1}^3 \delta^n a_n(Z_0, Z_1, Z_2) + \dots, \quad (3.2)$$

$$b = B_0 + \sum_{n=1}^3 \delta^n B_n(Z_0, Z_1, Z_2) + \dots, \quad (3.3)$$

$$\psi = \psi_0 + \sum_{n=1}^3 \delta^n \psi_n(Z_0, Z_1, Z_2) + \dots. \quad (3.4)$$

Using the chain rule, the spatial derivative becomes

$$\frac{d}{dZ} = D_0 + \delta D_1 + \delta^2 D_2 + \dots, \quad (3.5)$$

where $D_n = \partial/\partial Z_n$. The delay parameter ϵ is ordered as

$$\epsilon = \epsilon_0 + \delta^2 \epsilon_2, \quad (3.6)$$

where ϵ_0 is the critical value such that (2.10) is not satisfied, (i.e. ϵ_0 is a solution of (2.12)). This is standard for this method [11], as it allows the influence from the nonlinear terms and the control parameter to occur at the same order.

Using (3.2)–(3.6) in (2.4a)–(2.4c) and equating like powers of δ yields equations at $O(\delta^i)$, $i = 1, 2, 3$ of the form:

$$L_1(a_i, B_i, \psi_i) = S_{i,1}, \quad (3.7)$$

$$L_2(a_i, B_i, \psi_i) = S_{i,2}, \quad (3.8)$$

$$L_3(a_i, B_i, \psi_i) = S_{i,3}, \quad (3.9)$$

where, the L_i , $i = 1, 2, 3$ are the differential operators

$$L_1(a_i, B_i, \psi_i) = D_0 a_i - B_i \equiv S_{i,1}, \quad (3.10)$$

$$\begin{aligned} L_2(a_i, B_i, \psi_i) &= D_0 B_i - \psi_0^2 a_i - 2a_0 \psi_0 \psi_i \\ &+ \gamma_1 \{ \gamma_{20} a_i + v [b_1 B_i + c_1 (\psi_0 a_i + a_0 \psi_i)] \\ &- 3\gamma_{30} a_0^2 a_i - 5\gamma_{40} a_0^4 a_i \} \equiv S_{i,2}, \end{aligned} \quad (3.11)$$

$$\begin{aligned} L_3(a_i, B_i, \psi_i) &= a_0 (D_0 \psi_i) + 2(\psi_0 B_i + B_0 \psi_i) \\ &- \gamma_1 \{ \gamma_{50} a_i + v [c_1 B_i - b_1 (\psi_0 a_i + a_0 \psi_i)] \\ &- 3\gamma_{60} a_0^2 a_i - 5\gamma_{70} a_0^4 a_i \} \equiv S_{i,3}, \end{aligned} \quad (3.12)$$

where $\gamma_p = \gamma_{p0} + \delta^2 \gamma_{p2}$ with $p = 2, 3, \dots, 7$, the source terms $S_{i,j}$ for $i, j = 1, 2, 3$ at $O(\delta)$, $O(\delta^2)$, and $O(\delta^3)$ are given by the following:

$O(\delta)$:

$$S_{1,1} = 0 \quad (3.13)$$

$$S_{1,2} = 0 \quad (3.14)$$

$$S_{1,3} = 0. \quad (3.15)$$

$O(\delta^2)$:

$$S_{2,1} = -D_1 a_1 \quad (3.16)$$

$$S_{2,2} = -D_1 B_1 + a_0 \psi_1^2 + 2\psi_0 a_1 \psi_1 - \gamma_1(\gamma_{22} a_0 + v c_1 \psi_1 a_1 - 3\gamma_{30} a_0 a_1^2 - \gamma_{32} a_0^3 - 10\gamma_{40} a_0^3 a_1^2 - \gamma_{42} a_0^5) \quad (3.17)$$

$$S_{2,3} = -a_0 D_1 \psi_1 - a_1 D_0 \psi_1 - 2\psi_1 B_1 + \gamma_1[(\gamma_{52} a_0 - v b_1 \psi_1 a_1) - (3\gamma_{60} a_0 a_1^2 + \gamma_{62} a_0^3) - (10\gamma_{70} a_0^3 a_1^2 + \gamma_{72} a_0^5)]. \quad (3.18)$$

$O(\delta^3)$:

$$S_{3,1} = -D_1 a_2 - D_2 a_1 \quad (3.19)$$

$$S_{3,2} = -D_1 B_2 - D_2 B_1 + 2a_2 \psi_0 \psi_1 + a_1(2\psi_0 \psi_2 + \psi_1^2) + 2a_0 \psi_1 \psi_2 - \gamma_1\{\gamma_{22} a_1 + v c_1(\psi_1 a_2 + \psi_2 a_1) - [\gamma_{30}(a_1^3 + 6a_0 a_1 a_2) + 3\gamma_{32} a_0^2 a_1] - [\gamma_{40}(10a_0^2 a_1^3 + 20a_0^3 a_1 a_2) + 5\gamma_{42} a_0^4 a_1]\} \quad (3.20)$$

$$S_{3,3} = -D_1 \psi_2 - D_2 \psi_1 - a_1(D_1 \psi_1 + D_0 \psi_2) - a_2 D_0 \psi_1 - 2(\psi_1 B_2 + \psi_2 B_1) + \gamma_1\{\gamma_{52} a_1 - v b_1(\psi_1 a_2 + \psi_2 a_1) - [\gamma_{60}(a_1^3 + 6a_0 a_1 a_2) + 3\gamma_{62} a_0^2 a_1] - [\gamma_{70}(10a_0^2 a_1^3 + 20a_0^3 a_1 a_2) + 5\gamma_{72} a_0^4 a_1]\}. \quad (3.21)$$

Also, (3.7) may be solved for B_i in terms of a_i . and ψ_i . Using this in (3.8) yields ψ_i

$$\psi_i = \frac{\theta_i}{\phi_1}, \quad (3.22)$$

where

$$\theta_i = -D_0 S_{i,1} + D_0^2 a_i - \psi_0^2 + \gamma_1\{\gamma_{20} a_i - 3\gamma_{30} a_0^2 a_i - 5\gamma_{40} a_0^4 a_i + v[b_1(-S_{i,1} + D_0 a_i) + c_1]\} - S_{i,2} \quad (3.23)$$

and

$$\phi_1 = 2a_0 \psi_0 - v \gamma_1 c_1 a_0. \quad (3.24)$$

Using (3.22) and the equation for B_i in (3.10) yields the composite equation:

$$L_c a_i \equiv \Gamma_i, \quad (3.25)$$

where

$$\Gamma_i \equiv S_{i,3} - \frac{a_0}{\phi_1} \left(D_0 \zeta_i \right) - \frac{2B_0}{\phi_1} \zeta_i - \gamma_1 v b_1 a_0 \frac{\zeta_i}{\phi_1} + (2\psi_0 - \gamma_1 v c_1) S_{i,1}, \quad (3.26)$$

$$\zeta_i = -D_0 S_{i,1} - \gamma_1 v b_1 S_{i,1} - S_{i,2}. \quad (3.27)$$

We shall now use (3.26) and (3.27) to systematically identify and suppress secular terms in the solutions of (3.10),(3.11),(3.12). Let us now turn to finding the solutions of (3.10),(3.11),(3.12). In what follows, we shall detail the solution of the above system of equations for the case $\epsilon_0 = \epsilon_{01}$. In order to achieve that we must find first the fixed points. The characteristic polynomial of the Jacobian matrix of a fixed point of (2.4a),(2.4b),(2.4c) may be expressed as

$$\lambda^3 + \delta_1 \lambda^2 + \delta_2 \lambda + \delta_3 = 0, \quad (3.28)$$

as in (2.8), and the fixed point values (a_i, ψ_i) are to be substituted in terms of the system parameters.

The condition $\delta_1 \delta_2 - \delta_3 = 0$ yields an involved equation in ϵ which actually can be solved easily numerically for ϵ_0 by the root method.

For $O(\delta)$ the Eqns. (3.13)–(3.15) give $S_{i,1} = S_{i,2} = S_{i,3} = 0$, and hence we may pick a solution for the first order as

$$a_1 = \alpha(Z_1, Z_2) e^{\lambda_1 Z_0} + \beta(Z_1, Z_2) e^{\lambda_2 Z_0} + \gamma(Z_1, Z_2) e^{\lambda_3 Z_0}, \quad (3.29)$$

where $\beta = \bar{\alpha}$ is the complex conjugate of α and $\lambda_2 = \lambda_1$. As evident for the Routh–Hurwitz condition, the α and β modes correspond to the center manifold where $\lambda_{1,2}$ are purely imaginary and where the Hopf bifurcation occurs, while γ corresponds to the attractive direction or the stable manifold. Since we wish to construct and analyze the stability of the periodic orbits which lie in the center manifold, we should take $\gamma = 0$ so (3.29) becomes

$$a_1 = \alpha(Z_1, Z_2) e^{i\omega Z_0} + \beta(Z_1, Z_2) e^{-i\omega Z_0}. \quad (3.30)$$

Using (3.13)–(3.15) for $i = 1$ then the first order fields (a_1, B_1, ψ_1) are

$$B_1 = D_0 a_1 = i\omega \alpha e^{i\omega Z_0} - i\omega \beta e^{-i\omega Z_0}, \quad (3.31)$$

and also (3.22) becomes

$$\begin{aligned} \psi_1 = & \frac{1}{\phi_1} \left[-\omega^2 - \psi_0^2 + \gamma_1 \left(\gamma_{20} + v c_1 \psi_0 - 3\gamma_{30} a_0^2 - 5\gamma_{40} a_0^4 \right) \right] \\ & \times \left(\alpha e^{i\omega Z_0} + \beta e^{-i\omega Z_0} \right) + \frac{\gamma_1 v b_1}{\phi_1} \left(i\omega \alpha e^{i\omega Z_0} - i\omega \beta e^{-i\omega Z_0} \right). \end{aligned} \quad (3.32)$$

Now that the first order solutions (3.30)–(3.32) are known, the second order sources S_{21} , S_{22} , S_{23} may be evaluated via (3.16)–(3.18). Using these sources in (3.26) we obtain Γ_2 which may be written as

$$\Gamma_2 = \Gamma_2^{(0)} + \Gamma_2^{(1)} e^{i\omega Z_0} + \Gamma_2^{(2)} e^{2i\omega Z_0} + c.c. \quad (3.33)$$

Setting the coefficients of the secular $e^{i\omega Z_0}$ terms (which are the solutions of the homogeneous equation for $i = 1$) to zero, i.e. $\Gamma_2^{(1)} = 0$ yields

$$\begin{aligned} D_1\alpha &= \frac{\partial\alpha}{\partial Z_1} = 0, \\ D_1\beta &= \frac{\partial\beta}{\partial Z_1} = 0. \end{aligned} \quad (3.34)$$

Using (3.34), the second order sources, and assuming a second-order particular solution for a_2 of the form:

$$a_2 = a_2^{(0)} + a_2^{(2)} e^{2i\omega Z_0}, \quad (3.35)$$

having the standard form of a time-independent or DC term plus second-harmonic terms, the composite equations (3.25)–(3.27) for $i = 2$, yield

$$L_c a_2 = \Gamma_2^{(0)} + \Gamma_2^{(2)} e^{2i\omega Z_0}, \quad (3.36)$$

which will be solved for the particular solution $a_2^{(0)}$, and $a_2^{(2)}$ by equating both sides of the expression (3.36). In terms of the operator L_c which is obtained from (3.26), the particular solution takes the form:

$$\begin{aligned} a_2^{(0)} &= -\Gamma_2^{(0)} [a_0(v c_1 \gamma_1 - 2\psi_0)] \left\{ 2B_0 [-\psi_0^2 + \gamma_1(\gamma_{20} - 3a_0^2\gamma_{30} - 5a_0^4\gamma_{40} + v c_1 \psi_0)] \right. \\ &\quad + a_0 \gamma_1 \{ (\gamma_{50} - 3a_0^2\gamma_{60} - 5a_0^4\gamma_{70})(v c_1 \gamma_1 - 2\psi_0) \\ &\quad \left. + v b_1 [\gamma_1(\gamma_{20} - 3a_0^2\gamma_{30} - 5a_0^4\gamma_{40}) + \psi_0^2] \} \right\}^{-1}, \end{aligned} \quad (3.37)$$

$$\begin{aligned} a_2^{(2)} &= -\Gamma_2^{(2)} [a_0(v c_1 \gamma_1 - 2\psi_0)] \left\{ 6a_0^2 B_0 \gamma_1 \gamma_{30} + 10a_0^4 B_0 \gamma_1 \gamma_{40} \right. \\ &\quad + 3a_0^3 \gamma_1 [\gamma_{30}(2i\omega + v b_1 \gamma_1) + \gamma_{60}(v c_1 \gamma_1 - 2\psi_0)] \\ &\quad + 5a_0^5 \gamma_1 [\gamma_{40}(2i\omega + v b_1 \gamma_1) + \gamma_{70}(v c_1 \gamma_1 - 2\psi_0)] \\ &\quad + 2B_0 [4\omega_2 - 2iv\omega b_1 \gamma_1 + \psi_0^2 - \gamma_1(\gamma_{20} + v c_1 \psi_0)] \\ &\quad + a_0 \{ 8i\omega^3 - 2iv^2\omega b_1^2 \gamma_1^2 - 6i\omega\psi_0^2 + 2\gamma_1\psi_0(3iv\omega c_1 + \gamma_{50}) \\ &\quad \left. + \gamma_1 [-2i\omega(v^2 c_1^2 \gamma_1 + \gamma_{20}) - v c_1 \gamma_1 \gamma_{50}] - v b_1 \gamma_1 (-8\omega_2 + \gamma_1 \gamma_{20} + \psi_0^2) \} \right\}^{-1}. \end{aligned} \quad (3.38)$$

Using (3.22), the second order sources and the equation for B_i in (3.10) with $i = 2$ then we can find the second order fields B_2 and ψ_2 . Substituting them into the (3.19)–(3.21) we find the third order sources and we may evaluate the coefficients of the secular term $e^{i\omega t}$ in the composite source Γ_3 of (3.26). Suppressing again the secular terms to obtain uniform expansions yields the final equation for the evolution of the coefficients in the linear solutions (3.30)–(3.32) on the slow second-order time scales

$$\frac{\partial\alpha}{\partial Z_2} = S_{1r}\alpha^2\beta + S_{2r}\alpha. \quad (3.39)$$

Writing $\alpha = \frac{1}{2}Ae^{i\theta}$ and separating (3.39) into real and imaginary parts, yields

$$\frac{\partial A}{\partial Z_2} = \frac{S_{1r}A^3}{4} + S_{2r}A, \quad (3.40)$$

where S_{1r} and S_{2r} represent the real parts of S_1 and S_2 respectively. In the usual way, the fixed points of (3.40), $(A_1, A_{2,3})$ where

$$\begin{aligned} A_1 &= 0, \\ A_{2,3} &= \pm 2\sqrt{-\frac{S_{2r}}{S_{1r}}} \end{aligned} \quad (3.41)$$

give the amplitude of the solution $\alpha = \frac{1}{2}Ae^{i\theta}$, with $A_{2,3}$ corresponding to the bifurcation periodic orbits. Clearly $A_{2,3}$ are real fixed points whenever

$$\frac{S_{2r}}{S_{1r}} < 0, \quad (3.42)$$

and the Jacobian of the right hand side of (3.40) evaluated at $A_{2,3}$ is $J|_{A_{2,3}} = -2S_{2r}$, where $J(A) = \frac{\partial(\frac{S_{1r}A^3}{4} + S_{2r}A)}{\partial A}$. Clearly, a necessary condition for stability is to have $S_{2r} > 0$, and for instability $S_{2r} < 0$. Thus, the system undergoes:

a. supercritical Hopf bifurcations when

$$S_{2r} > 0, \quad S_{1r} < 0, \quad (3.43)$$

b. subcritical Hopf bifurcations when

$$S_{2r} < 0, \quad S_{1r} > 0. \quad (3.44)$$

4 Discussion of Results

In this section, we consider the numerical results which follow from the analysis in the previous section. The fixed point equation (2.6) can be solved analytically for each fixed point x_i using the program Mathematica, for $i = 1, \dots, 4$. The characteristic polynomial of the Jacobian matrix of a fixed point of (2.4a), (2.4b), (2.4c) may be expressed as in (3.1). Since all coefficients α_i , for $i = 1 \dots 3$ depend on the nine system parameters, we fix $b_1 = 0.08$, $b_3 = -0.65$, $b_5 = 0.1$, $c_1 = 0.5$, $c_3 = 1$, $c_5 = -0.07$, $\omega = 0$, and $v = 0.01$. The possibility of bounded chaotic solitons depends on the system being fairly strongly dissipative near the fixed points $(a_0, 0, \psi_0)$ in a significant part of the phase space, with the strong dissipativity ruling out the appreciable volume expansion associated with an attractor at infinity, as well as volume-conserving quasiperiodic behavior. The trace of the Jacobian matrix for this sets of values at the fixed point $(a_0, 0, \psi_0)$, which gives the local logarithmic rate of change of (a, b, ψ) phase-space volume V is $\frac{1}{V} \frac{dV}{dt} = J(a_0, 0, \psi_0) = -0.0062$, where $J(a(z), b(z), \psi(z)) = \frac{\partial(a_z, b_z, \psi_z)}{\partial(a, b, \psi)}$, so we may anticipate that the orbits may go to an attractor at infinity, since the dissipation is weak.

The four fixed points can be analytically found as a function of only one parameter, in our case we chose ϵ as being the free parameter. By choosing “the right fixed point”, the Hopf curve $\alpha_1\alpha_2 - \alpha_3 = 0$ may be solved numerically for ϵ , which gives $\epsilon_0 = -0.0000807$. The idea is to find the “right” ϵ which will give rise to the condition for Hopf bifurcation, (i.e. $\alpha_1 > 0$, $\alpha_2 > 0$, $\alpha_3 > 0$ and $\alpha_1\alpha_2 - \alpha_3 < 0$).




Figure 1: Stable periodic oscillations on the limit cycle $a(z)$ vs. z




Figure 2: $b(z) = a_z$ vs. z

We obtain $\alpha_1 = 0.006$, $\alpha_2 = 0.001$, $\alpha_3 = 0.0001$ and $\alpha_1\alpha_2 - \alpha_3 = -1.01 \cdot 10^{-6}$ for an $\epsilon = -0.008$, i.e. $\epsilon < \epsilon_0$.

Now we will analyze the multiple scales method to construct the analytical approximations for the periodic orbits arising through the Hopf bifurcations of the fixed point. The delay parameter ϵ (or the bifurcation parameter) is ordered as $\epsilon = \epsilon_0 + \delta^2\epsilon_2$, where $\epsilon_0 = -0.0000807$, and $\epsilon_2 = -0.1$. This method allows the influence from the nonlinear terms and the control parameter to occur at the same order. For the system parameters chosen above, at the fixed points, we get $(a_0, 0, \psi_0) = (0.0121663, 0, -0.00514)$.

From (3.25) and by the method presented in Section 2, the final equation for the evolution coefficients in the linear solutions, on the slow second-order time scale is

$$\frac{\partial\alpha(Z_1, Z_2)}{\partial Z_2} = S_1\alpha^2(Z_1, Z_2)\beta(Z_1, Z_2) + S_2\alpha(Z_1, Z_2), \quad (4.1)$$

where $S_1 = -3235.55 + 295.279i$ and $S_2 = 297.074 - 32.26i$. Since $S_{2r} = \text{Re}(S_2) > 0$, and $S_{1r} = \text{Re}(S_1) < 0$, then this situation will correspond to a supercritical Hopf bifurcation.

Figures 1–3 show the time behaviors of $a(z)$, $b(z)$, and $\psi(z)$ for $\epsilon = -0.00008$ (the supercritical regime). Note that, as anticipated from before, there is a stable limit cycle attractor at ϵ , the solution remains positive and bounded while it stays periodic.




Figure 3: $\psi(z) = \phi_z$ vs. z

Clearly, similar stable periodic solutions may be obtained for many other parameter sets. For each case, the overall solution $A(x, t)$ of the CGLE is, via (2.13), a quasiperiodic solution.

One may also use the above approach to both explain, and extend, the numerical treatment by Bruschi et al [10, 9] of the periodic traveling waves of the CGLE using the bifurcation software AUTO [12]. However, the solutions in Bruschi et al do not appear to be clearly correlated to the dissipative solitons of the CGLE in Akhmediev et al [1]. Hence, we shall move on next to briefly consider possible generalizations of the above treatment.

5 Remarks on Generalized Hopf Bifurcations and Emergence of Global Structure

One may pursue the line of inquiry based on the traveling waves or spatial ODEs even further to track the emergence of global structure. We have done preliminary work along these lines. However, although there is a well-established roadmap and it has been implemented in detail for the well known Continuous Stirred Tank Reactor System [19, 30], we are not convinced of its relevance to the actual numerical simulations of dissipative solitons [1, 2]. Hence, we present it here as a possible future direction to pursue.

For completeness, let us first consider more degenerate cases where more than one root of the Jacobian is non-hyperbolic. In such cases the non-hyperbolic eigenvalues of the Jacobian matrix, may consist of either:

a. a double zero: $\lambda_{1,2} = 0 \quad \lambda_3 \in \mathfrak{R}$

b. one zero and a complex conjugate pair: $\lambda_1 = 0 \quad \lambda_2 = \bar{\lambda}_3$

c. a triple zero: $\lambda_{1,2,3} = 0$

For the above situations, we have the following sub-cases of the so-called “degenerate Hopf” ($H1$) bifurcation. Each sub-case is given a name:

$$F_1 : \quad \lambda_{1,2} = 0, 0 \tag{5.1}$$

$$F_2 : \quad \lambda_{1,2,3} = \pm i\omega_0, 0 \tag{5.2}$$

$$G_1 : \quad \lambda_{1,2,3} = 0, 0, 0 \quad (5.3)$$

In these cases, [30, 5, 20] and [24, 28, 29], these (*H1*) bifurcations may lead to global structure including homoclinic orbits, invariant tori, and period doubling to chaos at the (*H1*) points. One may also work perturbatively [5] near these (*H1*) points as done by Keener for the well-known Continuous Stirred Tank Reactor problem.

Two other degenerate/generalized Hopf bifurcation scenarios are possible. As seen in Chapter 3 (3.40), the normal form for the Hopf bifurcation may be written as

$$\dot{r} = r[\alpha(\mu) + c_1(\mu)r^2 + c_2(\mu)r^4 + \dots] \quad (5.4)$$

$$\dot{\theta} = \omega_0 + O(\mu, r^2) \quad (5.5)$$

where we have made the identification $A \rightarrow r$, $S_{1r}/4 \rightarrow c_1$, $S_{2r} \rightarrow \alpha$, and higher order nonlinear terms are included.

The first kind of possible degeneracy (the (*H2*) kind) occurs if

$$\begin{aligned} \alpha = \alpha' = \dots = \alpha^{(k)} &= 0 \\ \alpha^{(k+1)} &\neq 0. \end{aligned} \quad (5.6)$$

This is the so-called k^{th} order (*H2*) degeneracy and it gives rise to multiple Hopf points and multiple periodic orbits. The resulting structure is thus similar to that resulting from a regular Hopf bifurcation, and much less complex than the structure produced by (*H1*) bifurcation.

A second possible degeneracy in the normal form (5.4) corresponds to

$$\begin{aligned} c_1 = c_2 = \dots = c_m &= 0 \\ c_{m+1} &\neq 0. \end{aligned} \quad (5.7)$$

This so-called m^{th} order (*H3*) degeneracy results in isolated branches of periodic solutions unconnected to the main branch.

When the k^{th} order (*H2*) degeneracy and the m^{th} order (*H3*) degeneracy occur simultaneously, the normal form (5.4) may be rescaled to the form:

$$\dot{r} = r[r^{2m+2} + \dots \pm \mu^{k+1}] \quad (5.8)$$

This is the so-called H_{mk} degeneracy.

In the case of the (*H2*) degeneracy, the complex conjugate eigenvalues $\pm i\omega$ at the Hopf point cross the imaginary axis tangentially leading, after additional analysis, to multiple periodic orbits.

For (*H3*) degeneracy, one may obtain isolated branches (isolas) of periodic orbits unconnected to the main branch.

However, of greatest interest are the (*H1*) bifurcations where the Jacobian has more than one non-hyperbolic eigenvalue and global structure emerges. These will be pursued in future work, together with the (*H2*) and (*H3*) cases.

References

- [1] N. Akhmediev. Pulsating solitons in dissipative systems. 4th IMACS Conf. on Nonlinear Waves, Athens, Georgia, April 2005, 2005.
- [2] N. Akhmediev, J. Soto-Crespo, and G. Town. Pulsating solitons, chaotic solitons, period doubling, and pulse coexistence in mode-locked laser: CGLE approach. *Phys. Rev.E*, 63:56602, 2001.
- [3] R. Alvarez, M. van Hecke, and W. van Saarloos. Sources and sinks separating domains of left- and right-traveling waves: Experiment versus amplitude equations. *Phys.Rev.E*, 56:R1306, 1997.
- [4] I.S. Aranson and L.Kramer. The World of the Complex Ginzburg-Landau Equation. *Rev. Mod. Phys.*, 74:99, 2002.
- [5] V.I. Arnold. Lectures on bifurcations in versal families. *Russian Math. Surveys*, 27:54, 1972.
- [6] D. Artigas, L. Torner, and N. Akhmediev. Robust heteroclinic cycles in the one-dimension CGLE. *Opt. Comm.*, 143:322, 1997.
- [7] N.J. Balmforth. Solitary waves and homoclinic orbits. *Ann. Rev. Fluid Mechanics*, 27:335, 1995.
- [8] C. Bowman and A.C. Newell. Natural Patterns and Wavelets. *Rev. Mod. Phys.*, 70:289, 1998.
- [9] L. Brusch, A. Torcini, and M. Bär. Nonlinear analysis of the Eckhaus Instability: Modulated amplitude waves and phase chaos. *Phys. D*, 160:127, 2001.
- [10] L. Brusch, A. Torcini, M. van Hecke, M. G. Zimmermann, and M. Bär. Modulated amplitude waves and defect formation in the one-dimensional complex Ginzburg-Landau equation. *Phys D.*, 160:127, 2001.
- [11] R.K. Dodd, J.C. Eilbeck, J.D. Gibbon, and H.C. Morris. *Solitons and Nonlinear Wave Equations*. Academic, London, 1982.
- [12] E. Doedel and J. Kern'vez. AUTO: Software for continuation problems in ordinary differential equations. California Institute of Technology, Pasadena, CA.,1986.
- [13] A. Doelman. Slow time-periodic solutions of the GL equation. *Phys.D*, 40:156, 1989.
- [14] A. Doelman. Periodic and quasiperiodic solutions of degenerate modulation equations. *Phys.D*, 53:249, 1991.
- [15] A. Doelman. Traveling waves in the complex GL equation. *J. Nonlin. Sci*, 3:225, 1993.
- [16] P. G. Drazin and R. S. Johnson. *Solitons: an Introduction*. Cambridge U. Press, Cambridge, 1989.

- [17] P.G. Drazin and W.H. Reid. *Hydrodynamic Stability*. Cambridge U. Press, Cambridge, 1981.
- [18] J. Duan and P. Holmes. Fronts, domain walls and pulses in a generalized GL equation. *Proc. Edin. Math. Soc.*, 38:77, 1995.
- [19] M. Golubitski and D.G. Schaffer. *Singularities and groups in Bifurcation Theory*, volume 51. Springer, New York, 1985.
- [20] J. Guckenheimer and P. Holmes. *Nonlinear Oscillations, Dynamical Systems, and Bifurcations of Vector Fields*, volume 42. Springer-Verlag, New York, 1983.
- [21] P. Holmes. Spatial structure of time periodic solutions of the GL equation. *Phys.D*, 23:84, 1986.
- [22] L.R. Keefe. Dynamics of perturbed wavetrain solutions to the Ginzburg-Landau equation. *Stud. Appl. Math.*, 73:91, 1985.
- [23] M.J. Landman. Solutions of the GL equation of interest in shear flow. *Stud. Appl. Math.*, 76:187, 1987.
- [24] W.F. Langford. Periodic and steady-state mode interactions lead to tori. *SIAM J. Appl. Math.*, 37:22, 1979.
- [25] J. D. Murray. *Mathematical Biology*. Springer-Verlag, Berlin, 1989.
- [26] P.K. Newton and L. Sirovich. Instabilities of the Ginzburg-Landau equation: Periodic solutions. *Quart. Appl. Math.*, XLIV:49, 1984.
- [27] P.K. Newton and L. Sirovich. Instabilities of the Ginzburg-Landau equation Pt.II: Secondary bifurcations. *Quart. Appl. Math.*, XLIV:367, 1986.
- [28] L.M. Pismen. Dynamics of lumped chemically reacting systems near singular bifurcations points. *Chem. Eng. Sci.*, 39:1063, 1984.
- [29] L.M. Pismen. Dynamics of lumped chemically reacting systems near singular bifurcations points-II. Almost Hamiltonian dynamics. *Chem. Eng. Sci.*, 40:905, 1985.
- [30] J. Planeaux. *Bifurcation Phenomena in CSTR Dynamics*. PhD thesis, University of Minnesota, 1993.
- [31] J. Satsuma and N. Yajima. Initial-value problems of one-dimensional self-modulation of nonlinear-waves in dispersive media. *Progr. Theoret. Phys. Suppl.*, 55:284, 1974.
- [32] J. M. Soto-Crespo, N. Akhmediev, and G. Town. Interrelation between various branches of stable solitons in dissipative systems. *Optics Comm.*, 199:283, 2001.
- [33] M. van Hecke, C. Storm, and W. van Saarloos. Sources, sinks and wavenumber selection in coupled CGL equations. *Phys.D*, 134:1, 1999.
- [34] W. van Saarloos and P. C. Hohenberg. Fronts, pulses, sources and sinks in generalized complex Ginzburg-Landau equation. *Phys.D*, 56:303, 1992.